

Interactions between plant proteins/enzymes and other food components, and their effects on food quality

Chenyan Lv, Guanghua Zhao & Yong Ning

To cite this article: Chenyan Lv, Guanghua Zhao & Yong Ning (2017) Interactions between plant proteins/enzymes and other food components, and their effects on food quality, Critical Reviews in Food Science and Nutrition, 57:8, 1718-1728, DOI: [10.1080/10408398.2015.1023762](https://doi.org/10.1080/10408398.2015.1023762)

To link to this article: <https://doi.org/10.1080/10408398.2015.1023762>



Accepted author version posted online: 20 Jul 2015.
Published online: 20 Jul 2015.



[Submit your article to this journal](#)



Article views: 365



[View related articles](#)



[View Crossmark data](#)



Citing articles: 2 [View citing articles](#)

Interactions between plant proteins/enzymes and other food components, and their effects on food quality

Chenyan Lv^{a,b}, Guanghua Zhao^{a,c}, and Yong Ning^c

^aBeijing Advanced Innovation Center for Food Nutrition and Human Health, College of Food Science and Nutritional Engineering, China Agricultural University, Beijing, China; ^bKey Laboratory of Functional Dairy, Ministry of Education, Beijing, China; ^cSchool of Laboratory Medicine, Hubei University of Chinese Medicine, Wuhan, China

ABSTRACT

Plant proteins are the main sources of dietary protein for humans, especially for vegetarians. There are a variety of components with different properties coexisting in foodstuffs, so the interactions between these components are inevitable to occur, thereby affecting food quality. Among these interactions, the interplay between plant proteins/enzymes from fruits and vegetables, cereals, and legumes and other molecules plays an important role in food quality, which recently has gained a particular scientific interest. Such interactions not only affect the appearances of fruits and vegetables and the functionality of cereal products but also the nutritive properties of plant foods. Non-covalent forces, such as hydrogen bond, hydrophobic interaction, electrostatic interaction, and van der Waals forces, are mainly responsible for these interactions. Future outlook is highlighted with aim to suggest a research line to be followed in further studies.

KEYWORDS

Interactions; food quality; plant protein; binding; enzymes

Introduction

Proteins in various foodstuffs are the main source of essential amino acids, which are used by cells to build new proteins for different purposes. Human beings use proteins for growth, and to build hormones, antibodies, and enzymes that regulate chemical reactions in the body. Proteins are widely distributed in various foods, such as seafood, meats, dairy, eggs, fruits, vegetables, whole grains, legumes, and nuts. These proteins are major components in foods, which are mainly derived from plant and animal sources.

On the other hand, food is a complex system where many other components such as polyphenols (Deng et al., 2009; Li et al., 2012), polysaccharides (Wang et al., 2002), lipids, and metal ions (Deng et al., 2009) coexist with proteins/enzymes, and thus their interactions occur inevitably. As we know, food quality encompasses not only sensory properties (appearance, texture, taste, and aroma) but also nutritive values, chemical constituents, mechanical, and functional properties. The above-mentioned interactions between proteins and other food components may affect both sensory and nutritive properties of foodstuffs. Of these interactions, some are undesirable. For example, polyphenol oxidases (PPO) can form dark spots in fruits and vegetables by enzymatic browning (Robinson, 1991). In contrast, there are a few examples, such as palm dates, prunes, raisins, and black tea, in which browning is considered as a desirable process, as it increases the product quality (Tomás-Barberán and Espin, 2001). Therefore, to control such interactions during food processing appears to be crucial for

food quality. Prior to controlling these interactions, it is important to understand the mechanism by which proteins interact with other molecules in foods.

In addition, many proteins have been used extensively in food industry. Interactions between added proteins and small molecules also occur. Soybean proteins are included in a wide variety of formulated foods, and are developed as a main source of protein for vegetarians nowadays. Interactions between soybean protein and lipids can improve emulsification property by forming protein–lipid complex. Similarly, ferritin with high bioavailability is being developed as an alternative iron supplement now (Zhao, 2010; Liao et al., 2014). In addition, polyphenols in plant food have been identified to interact with ferritin through non-covalent bonds, which can improve the digestive stability of ferritin (Li et al., 2012; Wang et al., 2014). This review focuses on a recent progress in the interactions between plant protein and other components of food, and their effects on food quality. In addition, attention is also drawn on the possible impacts of food processing on these interactions.

Proteins in cereals

Cereal grains are grown in greater quantities and provide more food energy worldwide than any other type of crop. So far, cereals have become an important part of diet of many people. They include maize, sorghum, millets, wheat, rice, barley, oats, teff, and quinoa. Proteins from cereals, such as gluten protein, are usually developed as food additives due to their important

function in food emulsification or texturing. Therefore, the interactions between cereal proteins and other food components may affect the sensory perception and consumer acceptance of cereal food products.

Gluten protein

Gluten properties are largely responsible for the end-use quality of wheat in many food products. It has been proved that gluten has the main contribution to wheat dough properties with a predominant role determining both dough machinability and textural characteristics of food product. Gluten proteins comprise two main sub-fractions. One is glutenin that confers strength and elasticity to dough, and another is gliadin that impairs viscous properties to gluten dough (Khatkar et al., 1995). In order to obtain food products with high quality, numerous additives are employed in bakery by reinforcing the role of gluten, or acting as emulsifiers. These components may affect gluten properties by interacting with gluten proteins.

The major non-starch polysaccharides of wheat flour are pentosans. Pentosans originate from the endosperm, the cell wall, and the bran of wheat grain (Wang et al., 2002). Ability to immobilize water and to form viscous solutions or gels by covalent cross-linking are important attributes that can have direct functional implications on gluten formation and properties. Water extractable pentosans (WEP) interfere with gluten formation in both direct and indirect ways. On the one hand, WEP can compete with gluten for water during the first stage of dough formation, resulting in delay in the development time of gluten (Labat et al., 2002). This corresponds to the indirect effect of WEP on gluten formation. On the other hand, WEP are able to directly cross-link with gluten, consequently affecting the extensibility of dough and gluten (Wang et al., 2002). In addition, the ferulic acid in WEP is proposed to involve in the interaction of pentosans with gluten (Wang et al., 2004). Similarly, hydroxypropylmethylcellulose (HPMC) has been shown to interact with gluten protein as well. The presence of HPMC did not modify the viscoelastic behavior of gluten dough during cooling at 25°C. However, the presence of HPMC increased the

solubility of gluten proteins in sodium dodecyl sulfate; this is possibly because HPMC interferes with protein association and its further aggregation during heating by occupying the space of proteins in the gluten network (Rosell and Foegeding, 2007).

The effects of hydrocolloids on the functional properties of wheat bread have been investigated recently. κ -Carrageenan, sodium alginate, xanthan gum, pectin, and some cellulose derivatives can affect dough rheology, bread volume, crumb texture, and shelf life during storage to different extents by forming hydrophilic complexes with gluten proteins (Mandala and Sotirakoglou, 2005; Qiu et al., 2015). In detail, pectin and λ -carrageenan strengthened wheat dough, and sodium alginate augmented the extensibility of dough. The formed weak complexes are mainly either due to attraction between local dipoles of carbohydrate residues of polysaccharide and charged groups of protein or the formation of unstable Schiff bases between the aldehyde groups of polysaccharide and the ε -amino groups of proteins (Rosell and Foegeding, 2007). Similarly, emulsifiers, such as sodium stearyl lactylate (SSL), used widely in bakery product can induce protein folding, including an increase in α -helix conformation and a decrease in β -sheet, turns, and random coil (Fig. 1). The conformation change may be the result of low burial of tryptophan residues to a more hydrophobic environment and the low percentage area of C–H stretching band for GS 0.25 (Gluten + 0.25% SSL) (Gómez et al., 2013).

In addition, bread making starts by adding water (and other ingredients) to flour and applying kinetic energy (by mixing), thereby forming extensible dough that contains a developed gluten network. During mixing, dough entraps air. Wheat flour-free lipids are bounded or trapped within the gluten fraction and can align at the interface of gas cells (Pareyt et al., 2011). Research on wheat flour lipids demonstrated that lipids played a significant role during dough mixing, fermentation and proofing, baking, and bread storage (Chin et al., 2010). Meanwhile, bread loaf volume is sensitive to the composition, extractability, and overall content of wheat flour lipids. In consequence, the food quality, especially that of bakery products, can be significantly changed by other components in wheat flour, and such changes can help improve the properties of




Figure 1. (A) Structure of sodium stearyl lactylate (SSL); (B) scanning electron microscopy of gluten prepared with emulsifier (1.0%). Native gluten (G), gluten–SSL (GS1). Magnification: 5000 \times (Gómez et al., 2013).

gluten protein, especially emulsification property, during food processing.

Sorghum protein

Sorghum (*Sorghum bicolor* L. Moench) is an important food cereal in many parts of Africa, Asia, and the semi-arid tropics worldwide, and acts as a principal source of protein, vitamins, and minerals for millions of poor people living in these regions. Porridges appear to be the most common types of food prepared from sorghum by cooking with boiling water. Protein fractionation studies have shown that prolamins and glutelin are the principal protein fractions (Duodu et al., 2003). However, the nutritive value of sorghum grain is relatively low due to its resistance to protease digestion. Factors affecting wet cooked sorghum protein digestibility may be categorized into two main groups: endogenous factors (disulfide and non-disulfide cross-linking, kafirin hydrophobicity, and changes in protein secondary structure) and exogenous factors (grain organizational structure, polyphenols, phytic acid, and starch and non-starch polysaccharides) (Duodu et al., 2003). Previous studies showed that cooking of sorghum in the presence of β -mercaptoethanol increased protein digestibility by reducing covalent cross-linking (disulfide bond) with other amino acids in the same or another protein molecule (Hamaker et al., 1987).

One important characteristic of sorghum is the abundance of tannins. It has been established that proteins generally interact with tannins by means of hydrogen bonding, hydrophobic interaction, electrostatic attraction, and covalent bonding associated with oxidation (Butler et al., 1984). Such interaction may lead to precipitation because of the large size of tannins, rendering most of the proteins insoluble. As expected, more tannins (2–4%) in sorghum were found to strongly bind with sorghum proteins, which have a loose and open structure, and are rich in proline (Butler et al., 1984). However, the anti-nutritional effects of sorghum tannin may be alleviated by treating the grain with dilute aqueous ammonia, strong alkalis, and formaldehyde, or by dehulling (McGrath et al., 1982). Moreover, the production of tannin-free sorghum by genetic modification can also improve the nutritional quality of sorghum products.

Similarly, phytic acid with high concentration naturally occurs in the germ of sorghum. Phytic acid is highly charged with six phosphate groups, and forms insoluble complexes with proteins by interactions (Ryden and Selvendran, 1993). This leads to reduced protein digestibility, which was attributed to the possible formation of a phytate–protein complex, and the protein was less susceptible to enzymatic attack (Duodu et al., 2003). However, there was no significant correlation between the percentage improvement in protein digestibility and dietary total phytic acid concentration. Many questions remain unanswered regarding the effect of phytate on sorghum proteins in particular.

Sunflower protein

Sunflower (*Helianthus annuus* L.) is one of the larger sources of vegetable oil and protein of good nutritional quality. Sunflower flours and protein concentrates have potential food uses because of their high protein content, white color, bland flavor,

and absence of anti-nutritive factors (Robertson and Morrison, 1977). Sunflower seeds also contain significant quantities of phenolic compounds, which remain in flour after oil extraction. Phenolic compounds naturally present in sunflower seeds mainly are chlorogenic and caffeic acids. To improve the quality of sunflower protein meal, attempts have been made to remove these polyphenolic substances, for instance acidic butanol extraction, to obtain colorless sunflower isolates from sunflower meal (Prasad, 1988). The acidic butanol extraction can remove 90% of phenolics from sunflower meal, resulting in a great reduction of color and change in flavor of sunflower meal. While it is possible to reduce the content of these compounds in protein products by modifying the extraction procedure (Salgado et al., 2011), it is impossible to eliminate them completely due to their strong interaction with proteins (Salgado et al., 2011). Under neutral and alkaline conditions, sunflower proteins develop dark green and brown colors due to bonding with oxidation products of polyphenolic compounds, especially chlorogenic acid (Prasad, 1988). Nevertheless, their final color tone was more dependent on the conditions used in the preparation process than on the amount of phenolic compounds in the product. It has been reported that the hydrogen bond between the hydroxyl groups of phenolic compounds and the peptide bond in proteins is unusually strong. In aqueous solutions, such strong interaction by the hydrogen bond greatly favors the formation of complexes between phenols and proteins. On the other hand, the interaction between sunflower protein and phenolic compounds conferred the antioxidant properties on sunflower protein films. Consequently, these specific protein films in packaging are of potential usefulness for preserving oxidation-sensitive products (Salgado et al., 2012).

Proteins in legumes

Legumes are widely recognized as important sources of food and feed proteins. In many regions of the world, legume seeds are the unique supply of proteins in diet. Proteins are a major component of legume seeds. Their nutritional and functional properties dramatically affect the overall quality of seeds and their technological performances. Soybean is the most important member of legumes family in the world. In developed countries, proteins from soybean seeds are now regarded as versatile functional ingredients or biologically active components more than as essential nutrients. Moreover, other components in soybean seeds, such as flavones, polyphenols, and other food additives, may interact with soybean proteins, affecting their functional properties.

Soybean protein isolate

Soybean protein isolates (SPI) because of their desirable functional properties, high nutritional value, and associated health effects, have been employed in a variety of formulated foods, and are developed as a main source of protein for vegetarians, especially for Asian populations. The major protein components of soybean are glycinin and β -conglycinin, which are used as emulsifiers due to the surface active properties of their constitutive proteins in bakery products, chocolate, instant

products (milk powder), margarines, and mayonnaise (Rydhaag and Wilton, 1981).

However, other molecules such as lipids and isoflavones coexist with soybean protein in soybean products, thereby affecting the properties of soybean protein by interactions during food processing. It has been reported that interactions between lecithin and soy protein can enhance the emulsification activity of soybean protein by forming the protein–lipid complex (Beckwith, 1984), which may be attributed to the components, such as proteins and phospholipids, possessing charges or having the capability to be ionized in food emulsions. Meanwhile, pH has been identified as an important factor in the emulsifying activity of soy protein and lecithin (Fig. 2). Moreover, different behaviors displayed by soy protein isolates are due to different protein structure and pH values. In other words, the presence of lecithin can enhance the initial characteristics of emulsions and diminish the creaming rate in both systems (Comas et al., 2006).

Isoflavones have been postulated as responsible for at least a part of beneficial health effects of soybean consumption. The affinity between soybean proteins and isoflavones depended on their diverse polarity and hydrophobicity as well as their abilities to form hydrogen bonds, which may affect the emulsification activity of soybean protein. Furthermore, it has been reported that enthalpic interactions (such as hydrogen bonding) between genistin and proteins would appear to come into play at pH 3.5, 4.5, or 5.6, with the resulting affinities being weaker with β -conglycinin than with glycinin. Meanwhile, malonylgenistin would likewise undergo an enthalpic interaction with proteins at pH 4.5 and 5.6, whereas at pH 3.5 hydrophobic bonds are favored (Speroni et al., 2010). These results help us to select optimal conditions during food processing to get food products with good taste and appearance. Nevertheless, the effect of isoflavones on the structures and properties of soybean protein remains to be further determined.

Ferritin

Ferritin is abundant in legume seeds. Ferritin as an iron storage protein has been extensively studied recently (Harrison and Arosio, 1996; Zhao, 2010). From the standpoint of nutrition,

biofortification of staple food with iron caged within phytoferritin from legumes is believed to be an effective strategy to combat iron deficiency anemia, which affects ~ 2 billion people around the world. However, there are many other components coexisting with ferritin in foodstuffs, and thus their interactions could occur, resulting in a change in the property of ferritin. It has been identified that the reductants in foodstuff, such as anthocyanins, phenolic acids, and ascorbic acids, can induce iron release from ferritin cavity (Deng et al., 2009) without influencing the primary/secondary structure of ferritin. The iron release rate partially depends on the structures and chelating activities of reductants. For example, the order of iron release from soybean seed ferritin (SSF) is as follows: delphinidin > cyaniding > petunidin > malvidin > delphinidin-3-O-glucoside > petunidin-3-O-glucoside. More interestingly, pigments can inhibit ferritin degradation during iron release to different extents (Deng et al., 2009). Moreover, it has been reported that tannic acid and epigallocatechin gallate (EGCG) can induce ferritin association (Li et al., 2012; Wang et al., 2014) (Fig. 3). Hydrogen bond and hydrophobic interaction may be two main factors responsible for the interaction between tannic acid and EGCG. It was also found that ferritin association induced by these small molecules could further improve the digestive stability of ferritin in vitro, but the evidence in vivo has been lacking.

Another molecule that can induce iron release from ferritin is a reduced form of nicotinamide-adenine dinucleotide (NADH). This compound is also widely distributed in foodstuffs. However, the mechanism of iron release from ferritin induced by NADH is different from the molecules listed above. NADH cannot contact the iron core directly due to the larger size of NADH (1.5 nm) than the size of ferritin channels (Masuda et al., 2010). Instead, NADH molecules bind on the surface of ferritin shell close to the four-fold channel of pea seed ferritin (PSF), which is 1.58 nm from the tryptophan residues calculated by fluorescence resonance energy transfer (Lv et al., 2013; Fig. 4). The interaction between these has been ascribed to van der Waals interactions or hydrogen bonds, as suggested by isothermal titration calorimetry (ITC) measurement (Chaikuad et al., 2005). Furthermore, since plastid DNA coexists with ferritin in the amyloplast of legume seeds, their




Figure 2. Crystal structures of (A) glycinin, and (B) β -conglycinin.




Figure 3. (A) Chemical structure of anthocyanin; (B) crystal structure of phytoferritin; (C) the SSF aggregation was initiated by mixing SSF with different concentrations of tannic acid (6.8–6.8 $\mu\text{g/mL}$). Transmission electron micrographs of holoferritin in the (a) absence, and (b) presence of tannic acid (Li et al., 2012).

interactions have been also investigated recently. Results demonstrated that the presence of DNA can enhance the rate of protein association during iron uptake by ferritin (Yang et al., 2014). On the other hand, SSF exhibited a marked DNA-protective function against oxidative damage at a low loading of Fe^{2+} ($\leq 48 \text{ Fe}^{2+}/\text{shell}$) (Liao et al., 2012; Fig. 4). Thus, the interactions between ferritin and other molecules significantly affect the stability and iron content of ferritin.

Importantly, the existence of dietary factors, such as phytic acid, polyphenols, and calcium, may affect the ferritin iron absorption by humans. At cell levels, it has been suggested that tannic acid increased iron uptake from intact ferritin, possibly by interfering with ferritin or ferritin mineral core assembly

due to its amphoteric properties and releasing iron for absorption. However, other dietary factors, such as phytic acid, ascorbic acid, and calcium, have no effect on the ferritin iron absorption (Kalganekar and Lönnerdal, 2008). Therefore, the interaction between dietary factors and ferritin could improve the stability and iron bioavailability of ferritin, especially for the tannins, but research in vivo needs to be elucidated.

Proteins/enzymes in fruits and vegetables

One prominent feature of fruits and vegetables is that the proteins occurring in them mainly comprise enzymes. Although these enzymes are usually much lower in content as compared




Figure 4. (A) Crystal structure of four-fold channel of phytoferritin. The tryptophan residues are labeled with red color; the red circle indicated the putative location of NADH binding sites. (B) The DNA protective role of SSF upon aerobic addition of $48 \text{ Fe}^{2+}/\text{protein}$ to apoferritin in vitro. Lane 1: plasmid DNA; lane 2: plasmid DNA + $48\text{-}\mu\text{M FeSO}_4$; lane 3: plasmid DNA + $1\text{-}\mu\text{M apoSSF}$ + $48\text{-}\mu\text{M FeSO}_4$ (Liao et al., 2012).

to carbohydrate, they play a key role in the quality of fruits and vegetables, including appearance, aroma, flavor, hand-feel, mouth-feel, and chewing sounds. Consumers integrate all these sensory perceptions into the final judgment of acceptability of fruits and vegetables.

Fruits and vegetables usually have a very short post-harvest life because of their relatively high metabolic activity and high sensitivity to fungal attack. Furthermore, during handling, storage, and marketing, they are highly susceptible to physical damage, leading to disruption of their cellular structures and consequently a speedup of softening and browning phenomena (Chisari et al., 2007). Some of these damages are induced by the interactions between enzymes widely distributed in fruits and vegetables with small molecules (oxygen, metal ions, pigments, etc.). Phenolic compounds play an important role in the visual appearance of foods. Anthocyanin pigments are responsible for most of the blue, purple, red, and intermediate hues of plant-derived foods. Therefore, the interaction between phenolic compounds and plant PPOs is considered to have significant impact on plant food quality. Elucidating such interactions seems to be very important, and fortunately, great progress has been made in this area as bellow.

Polyphenol oxidases

Browning of damaged tissues of fresh fruits and vegetables mainly occurs from the oxidation of phenolic compounds, and contributes significantly to loss of quality. A large body of work reports the characterization of oxidative enzymes from various fruits and vegetables such as apples, grapes, pears, eggplants, and strawberries (Carbonaro and Mattera, 2001). The primary enzymes responsible for the browning reaction are PPOs. The PPOs usually have a dinuclear copper center which catalyzes to insert oxygen in an *ortho*- position to an existing hydroxyl group in an aromatic ring (Virador et al., 2010; Fig. 5). The structure of the active site of enzyme is highly conserved, in which copper is bound by six or seven histidine residues and a single cysteine residue (Mayer, 2006), and the presence of the seventh histidine unit binding copper contributes to high enzyme activity (Hernández-Romero1 et al., 2006).

In plants, the PPOs are predominantly located in chloroplast thylakoid membranes, and are thereby physically separated from its natural substrate phenolic compounds, which occur in vacuoles. However, upon any cell-damaging treatment, the

enzyme and substrates come into contact, leading to rapid oxidation of phenols (Chazarra et al., 2001). Chlorogenic acid, caffeic acid, epicatechin, and catechin are the polyphenols commonly found in fruits and vegetables, which can act as substrates for PPOs and be oxidized to quinones by oxygen in the presence of PPOs. In turn, such quinones are very reactive and can react with each other and other cellular components to generate a black or dark brown pigment called melanin. This causes dark spots to form in plant tissues, frequently leading to a decrease in the quality of fruits or vegetables, especially for fresh-cut ones. It has been proved that pH is crucial for the oxidation of polyphenols by PPO, not only due to the optimal condition for PPO activity but also to the ionization state of enzymes (Kazandjian and Klibanov, 1985). In addition, many inhibitors of PPO have been described, which have diverse chemical structures. For example, the inhibition of glucose and fructose showed that the increasing concentrations of sugar caused a progressive inactivation of both enzymes, and such inhibition was much more evident in strawberry PPO than in others (Chisari et al., 2007). Salicylic acid as another inhibitor has been proved to competitively inhibit the activity of PPO by forming hydrogen bond with amino acids in PPO (Zhou et al., 2015). Differently, the development of browning is desirable for improvement of product quality of plant foods (Tomás-Barberán and Espin, 2001). For example, black color due to enzymatic browning is considered a criterion of quality in certain dried products such as black tea, coffee, and prune skins.

Peroxidases

Peroxidase (POD) occurring in almost all vegetables is another oxido-reductase enzyme that plays a crucial role in enzymatic browning as well, since diphenols may function as reducing substrates in its reaction (Robinson, 1991). Peroxidases usually comprise a family of isozymes containing identical heme groups but differ in the precise composition of glycoprotein. Peroxidases normally increase in activity and number during ripening, and can combine with hydrogen peroxide to produce an activated complex that can react with a wide range of donor molecules (Reed, 1975), causing undesirable changes in food materials, including off-flavor, aroma, and color. The involvement of peroxidase in browning is reported by different research groups. So far, a number of peroxidases from different fruits and vegetables have been identified using SDS-PAGE




Figure 5. (A) The pathways for polyphenol oxidation by polyphenol oxidases (PPO); (B) crystal structure of polyphenol oxidases (Virador et al., 2010). The histidine residues were labeled with yellow color.

followed by specific activity staining (Préstamo and Manzano, 1993). Similarly, recent studies have shown that cantaloupe melon peroxidase activity appears to be consistent with that of ascorbate peroxidase (Lamikanra and Watson, 2000), and that peroxidase activity in minimally processed cantaloupe melons could be the result of a preservative response to increased oxidative stress in the cut fruit (Lamikanra and Watson, 2001).

To further understand the interactions between peroxidases and substrates, the crystal structure of horseradish peroxidase isozyme has been solved, and key residues (Phe residues, 142, 68, and 179) involved in direct interactions with aromatic donor molecules have been identified (Gajhede et al., 1997). In addition, ascorbic acid and other natural antioxidants have been shown to inhibit peroxidase activities. All antioxidants used are able to terminate oxidation by preventing the formation of free radicals (Hemeda and Klein, 1990; Lamikanra and Watson, 2001). It has been reported that temperature preconditioning treatment can suppress increase in the peroxidase activity of squash during subsequent storage at 5°C (Wang, 1995).

Importantly, peroxidase is considered as the most heat-stable enzyme in plants, and there is an empirical relationship between residual peroxidase activity and the development of off-flavors and off-odors in foods. Thermal inactivation kinetic studies in peroxidase (in the range of 70–100°C) exhibited biphasic curves, providing evidence for the presence of isoenzymes with different thermal stabilities (Morales-Blancas et al., 2002). Therefore, inadequate thermal processing can cause reactivation of peroxidase and a loss of food quality. The detailed mechanism behind the thermal inactivation remains unclear up to this time. In addition, a pH of 2.4 at 25°C with low chloride concentration causes total detachment of heme. Once the heme–protein interaction is disturbed, there is a loss of protein stability. It was concluded that lipid oxidative activity of peroxidase aggregates was either due to the increased heme exposure with a change in temperature or pH or the increased number of active sites induced by heme migration (Burnette, 1977). A better understanding of interactions between peroxidase and substrates should enable the production of improved human food products with improved flavor and overall quality, resulting in longer storage periods.

Phenylalanine ammonia-lyase

Phenylalanine ammonia-lyase (PAL, EC 4.3.1.5) was first discovered in barley seedlings by Koukol and Conn (1961), and it has become the most studied enzyme concerned with secondary metabolism in plants. PAL activity in plants increases in response to several kinds of stress, including wounding (Ke and Saltveit, 1989), exposure to ethylene, low temperature, and fungal infection. PAL as a key enzyme in phenolic synthesis catalyzes the first reaction in the biosynthesis of plant phenylpropanoid products (Fig. 6). The synthesized phenolic compounds can further be oxidized by PPO, producing brown polymers that contribute to tissue browning (Ke and Saltveit, 1989). In detail, PAL catalyzes the non-oxidative deamination of *L*-phenylalanine to form *trans*-cinnamic acid and a free ammonium ion, which can induce the biosynthesis of a large range of phenylpropanoid-derived secondary products, such as flavonoids and isoflavonoids, coumarins, lignins, wound-




Figure 6. (A) The pathway for phenolic synthesis catalyzed by phenylalanine ammonia-lyase (PAL); (B) The crystal structure of PAL (Calabrese et al., 2004).

protective hydroxycinnamic acid esters, and other phenolic compounds (Jones, 1984). High activity of PAL is associated with the accumulation of anthocyanins and other phenolic compounds in the fruit tissues of several species.

One of the major causes of quality loss in minimally processed lettuce (*Lactuca sativa* L.) is the browning of cut pieces induced by PAL activity (López-Gálvez et al., 1996). In addition, it has been reported that strawberry fruits have a development-dependent expression of PAL activity and accumulation of phenolic substances derived from the phenylpropanoid pathway (Cheng and Breen, 1991). Importantly, chitosan and chitin treatments can lead to increase in PAL activity, which has been demonstrated to be one of the earliest responses of plants to the onset of infection by pathogens, and are often considered as indicators of resistance (Khan et al., 2003). Chilling damage has also been shown to induce an increase in PAL activity (Lafuente et al., 2003). Accordingly, many distinctive developmental features of flesh fruits, such as a loss of astringency and appearance of characteristic color at ripening, are related to PAL activity and changes in the synthesis and accumulation of phenolic compounds.

Pectinases

Food texture is a major determinant of consumer acceptance and preference for fruits and vegetables (Van Buggenhout et al., 2009). Pectic substances account for about one-third of the dry substance of primary cell walls of fruits and vegetables. Previous studies on fruits and vegetables suggested that the interactions between the pectic substances and pectinases coexisting in fruits and vegetables might affect their quality (Kashyap et al., 2001). Pectinase is a complex macromolecule, which plays an important role in the mechanical properties of plant tissue and in (pre-)processed plant-based foods. They catalyze numerous pectin conversion reactions, strongly degrading the pectic substances, thereby impacting the quality of fruits,

vegetables, and the related intermediate and end products (Duvetter et al., 2009).

Generally, pectinesterases, known as pectinmethylhydrolase, are involved in changes in the pectic substances of fruits and vegetables during ripening, storing, and processing by catalyzing de-esterification of the methoxyl group of pectin. Meanwhile, depolymerizing enzymes can catalyze hydrolysis or cleaving of α -1, 4-glycosidic linkages between galacturonic monomers in pectic substances. In addition, protopectinases, protopectin-solubilizing enzymes, which liberate water-soluble and highly polymerized pectin from protopectin, have been reported to react with the polygalacturonic acid region of protopectin and with the polysaccharide chains that may connect polygalacturonic acid chain with cell wall constituents (Alkorta et al., 1998).

In food industry, pectinases are extensively used in increasing the yield of fruit and vegetable juices, controlling cloud stability in juices, enzymatic peeling of fruits, controlling the rheological properties of purees and pastes, engineering the texture of fruits and vegetables, manufacture of wine, extraction of pigments and food colorings, and so on. Commercial pectic enzymes are used in apple juice manufacturing to de-pectinize pressed juices to remove turbidity and prevent cloud-forming. Similar to other enzymes in fruits and vegetables, the pH and temperature may affect pectinase activity significantly (Ceci and Lozano, 1998). Therefore, controlling of interactions between pectinases and pectic substances during food storage and processing is of prime importance, since desirable or deleterious reactions can be tailored (accelerated or inhibited), meeting specific quality targets.

Effects of food processing on the interactions between proteins/enzymes and other food components

Food processing has been known to affect content, activity, and bioavailability of food components, and it also plays an important role in the interactions between plant protein and other food components. The traditional food processing methods include heat treatment, fermentation, and germination. Heat treatment has been usually used for the inactivation of enzymes in plant food such as fruits and vegetables (Morales-Blancas et al., 2002). Phytic acid of soy meal (SM) could influence protein and important mineral digestion of monogastric animals. Recent studies have demonstrated that two-stage temperature protocol achieves better phytic acid degradation during the solid-state fermentation of *A. oryzae*. Therefore, the fermented soy meal has lower anti-nutritional factors (phytic acid and oligosaccharides) and higher nutritional value for animal feed (Chen et al., 2014). As an interesting alternative to traditional food processing and preservation methods, high-pressure processing has a potential for food preservation because it can inactivate microorganisms and enzymes responsible for shortening the life of a product. In addition to lengthening the shelf life of food products, high hydrostatic pressure (HHP) can modify functional properties of components such as proteins, which in turn can lead to the development of new products (Hendrickx et al., 1998). Moreover, ultrasound has attracted considerable interest in food science and technology because of

its promising effects in food processing and preservation. As one of the advanced food technologies, it can be applied to develop gentle but targeted processes to improve the quality and safety of processed foods, and offers the potential for improving existing processes as well as developing new process options (Knorr et al., 2004).

High hydrostatic pressure has been shown to improve protein solubility and dispersion stability of mineral-added SPI. In detail, HHP-denatured soybean proteins may coexist with different minerals at different pH values in the form of soluble species (Manassero et al., 2015). Thermal- and HHP-denatured calcium-added soybean proteins exhibited different solubility values. HHP may lead to dissociation of calcium from binding sites of soybean proteins, whereas thermal treatment cannot do so. Similarly, high intensity ultrasonic (HUS) pre-treatment can affect the properties of soybean protein. It has been shown that the surface hydrophobicity and free sulfhydryl (SH) content of SPI can increase with HUS-treatment time (Hu et al., 2013), which will further affect interaction forces between proteins and small molecules.

As we all know, enzymes are responsible for the quality of fruits and vegetables. In order to improve the quality of fruits and vegetables, enzymes have been inactivated by many methods to prevent interactions between enzymes and their substrates. As for thermal technologies, in addition to traditional heating, there are several methods, such as ohmic heating, that can raise temperature to a critical level in a very short time (Jaeger et al., 2010). Although traditional heat treatments can ensure safety and extend the shelf life of juices, undesirable brown color develops as the result of Maillard reaction between amino and carbonyl compounds. In contrast, high-pressure (HP) processing (at low and moderate temperatures) has a limited effect on pigments (chlorophyll, carotenoids, and anthocyanins) responsible for color and flavor in fruits and vegetables (Oey et al., 2008). However, due to cell disruption, high-pressure processing facilitates the occurrence of enzymatic and non-enzymatic reactions related to the texture of fruits and vegetables. This is because substrates, ions, and enzymes located in different compartments of cells can be liberated to interact with each other during high-pressure treatment (Oey et al., 2008). Moreover, pressure can enhance pectinmethyl-esterase (PME) action and lower polygalacturonase (PG) activity (occurring mostly at moderate temperature). Pectinases from different sources (Van den Broeck et al., 2000; Ly Nguyen et al., 2002) exhibit differences in their pressure and temperature stability. Consequently, different pressure and temperature combinations can be used to activate or inactivate some specific pectinases during processing to create textures, which cannot be formed by thermal processing.

In order to avoid detrimental changes in sensory and nutritive properties, pulsed electric field (PEF) pasteurization of fruit juices is a promising preservation method (Jaeger et al., 2010). It has been reported that PEF-processed juices had a lower 5-hydroxymethylfurfural concentration than those treated with heat, a fact that can be attributed to the reduced thermal load to which the product is exposed during PEF preservation. Thus, it seems that a combination of non-thermal and thermal

technologies could improve food quality better during food processing.

Conclusions and perspectives

Interactions between proteins and other molecules in different food systems have been extensively investigated recently. The crucial enzymes (PPO, POD, or PAL) in fruits and vegetables responsible for enzymatic browning should be inactivated or activated according to the need of customer. Interactions between plant proteins and other molecules, such as polysaccharides, lipids, and metal ions, occur constantly during food processing. These studies mainly focus on the interactions in vitro, and their effects on protein structures and functions. Non-covalent forces such as hydrogen bond, hydrophobic interaction, electrostatic interaction, and van der Waals forces are responsible for these interactions. Meanwhile, the pH value of an aqueous solution is also crucial for interactions due to the electrical charge of the molecules involved. All these interactions can further affect the appearance, nutrition, and texture of food. What's more, food-processing technologies can significantly affect interactions between plant protein and other molecules. Therefore, during food processing, it is of utmost importance to select optimal conditions to control these interactions for different applications. It is likely that other unidentified conditions related to the interaction coexist, which also make important contributions to this process.

However, there have been some questions that remain to be answered. First, better evaluation systems should be established to assess the effect of interactions on food quality. Second, the mechanism of interactions between plant protein and other molecules needs more detailed information to better control such interactions. Third, most of the interactions studied focus mainly on bi-molecule system. However, interactions in a real food system are much more complicated than those in a bi-molecule system. Finally, the bioavailability of entire foodstuff, which is a major concern during food intake, has not yet been elucidated.

Funding

The work was supported by the National Natural Science Foundation of China (31471693).

References

- Alkorta, I., Garbisu, C., Llama, M. J., and Serra, J. L. (1998). Industrial applications of pectic enzymes: A review. *Process Biochem.* **33**:21–28.
- Beckwith, A. C. (1984). Interaction of phosphatidylcholine vesicles with soybean 7S and 11S globulin proteins. *J. Agri. Food Chem.* **32**:1397–1402.
- Burnette, F. S. (1977). Peroxidase and its relationship to food flavor and quality: A review. *J. Food Sci.* **42**:1–6.
- Butler, L. G., Riedl, D. J., Lebryk, D. G., and Blytt, H. J. (1984). Interaction of proteins with sorghum tannin: Mechanism, specificity and significance. *J. Am. Oil Chem. Soc.* **61**:916–920.
- Calabrese, J. C., Jordan, D. B., Boodhoo, A., Sariaslani, S., and Vannelli, T. (2004). Crystal structure of phenylalanine ammonia lyase: Multiple helix dipoles implicated in catalysis. *Biochemistry.* **43**:11403–11416.
- Carbonaro, M., and Mattera, M. (2001). Polyphenoloxidase activity and polyphenol levels in organically and conventionally grown peach (*Prunus persica* L., cv. Regina bianca) and pear (*Pyrus communis* L., cv. Williams). *Food Chem.* **72**:419–424.
- Ceci, L., and Lozano, J. (1998). Determination of enzymatic activities of commercial pectinases for the clarification of apple juice. *Food Chem.* **61**:237–241.
- Chaikuad, A., Fairweather, V., Connors, R., Joseph-Horne, T., Turgut-Balik, D., and Brady, R. L. (2005). Structure of lactate dehydrogenase from *Plasmodium vivax*: Complexes with NADH and APADH. *Biochemistry.* **44**:16221–16228.
- Chazarra, S., García-Carmona, F., and Cabanes, J. (2001). Evidence for a tetrameric form of iceberg lettuce (*Lactuca sativa* L.) polyphenol oxidase: Purification and characterization. *J. Agr. Food Chem.* **49**:4870–4875.
- Chen, L., Vadlani, P. V., and Madl, R. L. (2014). High-efficiency removal of phytic acid in soy meal using two-stage temperature-induced *Aspergillus oryzae* solid-state fermentation. *J. Sci. Food Agric.* **94**:113–118.
- Cheng, G. W., and Breen, P. J. (1991). Activity of phenylalanine ammonia-lyase (PAL) and concentrations of anthocyanins and phenolics in developing strawberry fruit. *J. Am. Soc. Hortic Sci.* **116**:865–869.
- Chin, N. L., Rahman, R. A., Hashim, D. M., and Kowng, S. Y. (2010). Palm oil shortening effects on baking performance of white bread. *J. Food Process Eng.* **33**:413–433.
- Chisari, M., Barbagallo, R. N., and Spagna, G. (2007). Characterization of polyphenol oxidase and peroxidase and influence on browning of cold stored strawberry fruit. *J. Agr. Food Chem.* **55**:3469–3476.
- Comas, D. I., Wagner, J. R., and Tomás, M. C. (2006). Creaming stability of oil in water (O/W) emulsions: Influence of pH on soybean protein–lecithin interaction. *Food Hydrocolloid.* **20**:990–996.
- Deng, J., Cheng, J., Liao, X., Zhang, T., Leng, X., and Zhao, G. (2009). Comparative study on iron release from soybean (*Glycine max*) seed ferritin induced by anthocyanins and ascorbate. *J. Agr. Food Chem.* **58**:635–641.
- Duodu, K. G., Taylor, J. R. N., Belton, P. S., and Hamaker, B. R. (2003). Factors affecting sorghum protein digestibility. *J. Cereal Sci.* **38**:117–131.
- Duvetter, T., Sila, D. N., Van Buggenhout, S., Jolie, R., Van Loey, A., and Hendrickx, M. (2009). Pectins in processed fruit and vegetables: Part I—stability and catalytic activity of pectinases. *Compr. Rev. Food Sci. F.* **8**:75–85.
- Gajhede, M., Schuller, D. J., Henriksen, A., Simth, A. T., and Poulos, T. L. (1997). Crystal structure of horseradish peroxidase C at 2.15 Å resolution. *Nat. Struct. Mol. Biol.* **4**:1032–1038.
- Gómez, A. V., Ferrer, E. G., Añón, M. C., and Puppo, M. C. (2013). Changes in secondary structure of gluten proteins due to emulsifiers. *J. Mol. Struct.* **1033**:51–58.
- Hamaker, B. R., Kirleis, A. W., Butler, L. G., Axtell, J. D., and Mertz, E. T. (1987). Improving the in vitro protein digestibility of sorghum with reducing agents. *P. Natl. Acad. Sci. USA.* **84**:626–628.
- Harrison, P. M., and Arosio, P. (1996). The ferritins: Molecular properties, iron storage function and cellular regulation. *Biochim. Biophys. Acta Bioenerg.* **1275**:161–203.
- Hemeda, H. M., and Klein, B. P. (1990). Effects of naturally occurring antioxidants on peroxidase activity of vegetable extracts. *J. Food Sci.* **55**:184–185.
- Hendrickx, M., Ludikhuyze, L., Van den Broeck, I., and Weemaes, C. (1998). Effects of high pressure on enzymes related to food quality. *Trends Food Sci. Tech.* **9**:197–203.
- Hernández-Romero, D., Sanchez-Amat, A., and Solano, F. (2006). A tyrosinase with an abnormally high tyrosine hydroxylase/dopa oxidase ratio. *FEBS J.* **273**:257–270.
- Hu, H., Li-Chan, E. C., Wan, L., Tian, M., and Pan, S. (2013). The effect of high intensity ultrasonic pre-treatment on the properties of soybean protein isolate gel induced by calcium sulfate. *Food Hydrocolloid.* **32**:303–311.
- Jaeger, H., Janositz, A., and Knorr, D. (2010). The Maillard reaction and its control during food processing. The potential of emerging technologies. *Pathol. Biol.* **58**:207–213.
- Jones, D. H. (1984). Phenylalanine ammonia-lyase: Regulation of its induction, and its role in plant development. *Phytochemistry.* **23**:1349–1359.

- Kalgaonkar, S., and Lönnerdal, B. (2008). Effects of dietary factors on iron uptake from ferritin by Caco-2 cells. *J. Nutr. Biochem.* **19**:33–39.
- Kashyap, D. R., Vohra, P. K., Chopra, S., and Tewari, R. (2001). Applications of pectinases in the commercial sector: A review. *Bioresour. Technol.* **77**:215–227.
- Kazandjian, R. Z., and Klivanov, A. M. (1985). Regioselective oxidation of phenols catalyzed by polyphenol oxidase in chloroform. *J. Am. Chem. Soc.* **107**:5448–5450.
- Ke, D., and Saltveit, M. E. (1989). Wound-induced ethylene production, phenolic metabolism and susceptibility to russet spotting in iceberg lettuce. *Physiol. Plantarum.* **76**:412–418.
- Khan, W., Prithiviraj, B., and Smith, D. L. (2003). Chitosan and chitin oligomers increase phenylalanine ammonia-lyase and tyrosine ammonia-lyase activities in soybean leaves. *J. Plant Physiol.* **160**:859–863.
- Khatkar, B. S., Bell, A. E., and Schofield, J. D. (1995). The dynamic rheological properties of glutes and gluten sub-fractions from wheats of good and poor bread making quality. *J. Cereal Sci.* **22**:29–44.
- Knorr, D., Zenker, M., Heinz, V., and Lee, D. U. (2004). Applications and potential of ultrasonics in food processing. *Trends Food Sci. Tech.* **15**:261–266.
- Koukol, J., and Conn, E. E. (1961). The metabolism of aromatic compounds in higher plants IV. Purification and properties of the phenylalanine deaminase of *Hordeum vulgare*. *J. Biol. Chem.* **236**:2692–2698.
- Labat, E., Rouau, X., and Morel, M. H. (2002). Effect of flour water-extractable pentosans on molecular associations in gluten during mixing. *LWT Food Sci. Technol.* **35**:185–189.
- Lafuente, M. T., Zacarias, L., Martínez-Téllez, M. A., Sanchez-Ballesta, M. T., and Granell, A. (2003). Phenylalanine ammonia-lyase and ethylene in relation to chilling injury as affected by fruit age in citrus. *Postharvest Biol. Technol.* **29**:309–318.
- Lamikanra, O., and Watson, M. A. (2000). Cantaloupe melon peroxidase: Characterization and effects of additives on activity. *Nahrung.* **44**:168–172.
- Lamikanra, O., and Watson, M. A. (2001). Effects of ascorbic acid on peroxidase and polyphenoloxidase activities in fresh-cut cantaloupe melon. *J. Food Sci.* **66**:1283–1286.
- Li, M., Jia, X., Yang, J., Deng, J., and Zhao, G. (2012). Effect of tannic acid on properties of soybean (*Glycine max*) seed ferritin: A model for interaction between naturally occurring components in foodstuffs. *Food Chem.* **133**:410–415.
- Liao, X., Lv, C., Zhang, X., Masuda, T., Li, M., and Zhao, G. (2012). A novel strategy of natural plant ferritin to protect DNA from oxidative damage during iron oxidation. *Free Radical Biol. Med.* **53**:375–382.
- Liao, X., Yun, S., and Zhao, G. (2014). Structure, function, and nutrition of phytoferritin: A newly functional factor for iron supplement. *Crit. Rev. Food Sci. Nutr.* **54**:1342–1352.
- López-Gálvez, G., Saltveit, M., and Cantwell, M. (1996). Wound-induced phenylalanine ammonia lyase activity: Factors affecting its induction and correlation with the quality of minimally processed lettuces. *Postharvest Biol. Tech.* **9**:223–233.
- Lv, C., Bai, Y., Yang, S., Zhao, G., and Chen, B. (2013). NADH induces iron release from pea seed ferritin: A model for interaction between coenzyme and protein components in foodstuffs. *Food Chem.* **141**:3851–3858.
- Ly Nguyen, B., Van Loey, A., Fachin, D., Verlent, I., Duvetter, T., Vu, T. S., et al. (2002). Strawberry pectin methylesterase: Purification, characterization, thermal and high-pressure inactivation. *Biotechnol. Progr.* **18**:1447–1450.
- Manassero, C., Sergio, R., Vaudagna, S. R., Anón, M. C., and Speroni, F. (2015). High hydrostatic pressure improves protein solubility and dispersion stability of mineral-added soybean protein isolate. *Food Hydrocolloid.* **43**:629–635.
- Mandala, I. G., and Sotirakoglou, K. (2005). Effect of frozen storage and microwave reheating on some physical attributes of fresh bread containing hydrocolloids. *Food Hydrocolloid.* **19**:709–719.
- Masuda, T., Goto, F., Yoshihara, T., and Mikami, B. (2010). Crystal structure of plant ferritin reveals a novel metal binding site that functions as a transit site for metal transfer in ferritin. *J. Biol. Chem.* **285**:4049–4059.
- Mayer, A. M. (2006). Polyphenol oxidases in plants and fungi: Going places? A review. *Phytochemistry.* **67**:2318–2331.
- McGrath, R. M., Kaluza, W. Z., Daiber, K. H., Van der Riet, W. B., and Glenne, C. W. (1982). Polyphenols of sorghum grain, their changes during malting and their inhibitory nature. *J. Agr. Food Chem.* **30**:450–456.
- Morales-Blancas, E. F., Chandia, V. E., and Cisneros-Zevallos, L. (2002). Thermal inactivation kinetics of peroxidase and lipoxygenase from broccoli, green asparagus and carrots. *J. Food Sci.* **67**:146–154.
- Pareyt, B., Finnie, S. M., Putseys, J. A., and Delcour, J. A. (2011). Lipids in bread making: Sources, interactions, and impact on bread quality. *J. Cereal Sci.* **54**:266–279.
- Prasad, D. T. (1988). Studies on the interaction of sunflower albumins with chlorogenic acid. *J. Agr. Food Chem.* **36**:450–452.
- Préstamo, G., and Manzano, P. (1993). Peroxidases of selected fruits and vegetables and the possible use of ascorbic acid as an antioxidant. *HortScience.* **28**:48–50.
- Oey, I., Lille, M., Van Loey, A., and Hendrickx, M. (2008). Effect of high-pressure processing on colour, texture and flavour of fruit- and vegetable-based food products: A review. *Trends Food Sci. Tech.* **19**:320–328.
- Qiu, C., Zhao, M., and McClements, D. J. (2015). Improving the stability of wheat protein-stabilized emulsions: Effect of pectin and xanthan gum addition. *Food Hydrocolloid.* **43**:377–387.
- Reed, G. (1975). Oridoreductases. In: *Enzymes in Food Processing*, pp 216. Academic Press, New York, NY.
- Robertson, J. A., and Morrison, W. H. (1977). Effect of heat and frying on sunflower, oil stability. *J. Am. Oil Chem. Soc.* **54**:A77–A81.
- Robinson, D. S. (1991). Peroxidase and catalase. In: *Oxidative Enzymes in Foods*, pp. 1–49. Robinson, D. S., and Eskin, N. A. M., Eds., Elsevier, Amsterdam, the Netherlands.
- Rosell, C. M., and Foegeding, A. (2007). Interaction of hydroxypropylmethylcellulose with gluten proteins: Small deformation properties during thermal treatment. *Food Hydrocolloid.* **21**:1092–1100.
- Ryden, P., and Selvendran, R.R. (1993). Phytic acid: Properties and determination. In: *Encyclopaedia of Food Science, Food Technology and Nutrition*, pp. 3582–3587. Macrae, R., Robinson, R. K., and Sadler, M. J., Eds., Academic Press, London.
- Rydhag, L., and Wilton, I. (1981). The function of phospholipids of soybean lecithin in emulsions. *J. Am. Oil Chem. Soc.* **58**:830–837.
- Salgado, P. R., López-Caballero, M. E., Gómez-Guillén, M. C., Mauri, A. N., and Montero, M. P. (2012). Exploration of the antioxidant and antimicrobial capacity of two sunflower protein concentrate films with naturally present phenolic compounds. *Food Hydrocolloid.* **29**:374–381.
- Salgado, P. R., Ortiz, S. E. M., Petruccielli, S., and Mauri, A. N. (2011). Sunflower protein concentrates and isolates prepared from oil cakes have high water solubility and antioxidant capacity. *J. Am. Oil Chem. Soc.* **88**:351–360.
- Speroni, F., Milesi, V., and Anón, M. C. (2010). Interactions between isoflavones and soybean proteins: Applications in soybean-protein-isolate production. *LWT Food Sci. Technol.* **43**:1265–1270.
- Tomás-Barberán, F. A., and Espin, J. C. (2001). Phenolic compounds and related enzymes as determinants of quality in fruits and vegetables. *J. Sci. Food Agr.* **81**:853–876.
- Van Buggenhout, S., Sila, D. N., Duvetter, T., Van Loey, A., and Hendrickx, M. (2009). Pectins in processed fruits and vegetables: Part III – Texture engineering. *Compr. Rev. Food Sci. F.* **8**:105–117.
- Van den Broeck, I., Ludikhuyze, L. R., Van Loey, A. M., and Hendrickx, M. E. (2000). Inactivation of orange pectinesterase by combined high-pressure and temperature treatments: A kinetic study. *J. Agr. Food Chem.* **48**:1960–1970.
- Virador, V. M., Reyes Grajeda, J. P., Blanco-Labra, A., Mendiola-Olaya, E., Smith, G. M., Moreno, A., and Whitaker, J. R. (2010). Cloning, sequencing, purification, and crystal structure of Grenache (*Vitis vinifera*) polyphenol oxidase. *J. Agr. Food Chem.* **58**:1189–1201.
- Wang, C. Y. (1995). Effect of temperature preconditioning on catalase, peroxidase, and superoxide dismutase in chilled zucchini squash. *Postharvest Biol. Tech.* **5**:67–76.

- Wang, M., Hamer, R. J., van Vliet, T., and Oudgenoeg, G. (2002). Interaction of water extractable pentosans with gluten protein: Effect on dough properties and gluten quality. *J. Cereal Sci.* **36**:25–37.
- Wang, M., van Vliet, T., and Hamer, R. J. (2004). How gluten properties are affected by pentosans. *J. Cereal Sci.* **39**:395–402.
- Wang, A., Zhou, K., Qi, X., and Zhao, G. (2014). Phytoferritin association induced by EGCG inhibits protein degradation by proteases. *Plant Foods Hum. Nutr.* **69**:386–391.
- Yang, R., Yang, S., Liao, X., Deng, J., and Zhao, G. (2014). The interaction of DNA with phytoferritin during iron oxidation. *Food Chem.* **153**:292–297.
- Zhao, G. (2010). Phytoferritin and its implications for human health and nutrition. *Biochim. Biophys. Acta.* **1800**:815–823.
- Zhou, D., Li, L., Wu, Y., Fan, J., and Ouyang, J. (2015). Salicylic acid inhibits enzymatic browning of fresh-cut Chinese chestnut (*Castanea Mollissima*) by competitively inhibiting polyphenol oxidase. *Food Chem.* **171**:19–25.