

## Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

### Substitution of Saturated Fat in Processed Meat Products: A Review

J. C. Ospina-E<sup>a</sup>, A. Sierra-C<sup>a</sup>, O. Ochoa<sup>a</sup>, J. A. Pérez-Álvarez<sup>b</sup> & J. Fernández-López<sup>b</sup>

<sup>a</sup> Industria de Alimentos Zenú S.A.S. Cra 64C N° 104-03. Medellín, Colombia Centro de Investigación y Desarrollo Cárnico CI+D

<sup>b</sup> Universidad Miguel Hernández de Elche. Crta. a Beniel. Km. 3,2. Orihuela, Alicante, España. Grupo REVIV, Generalitat Valenciana y Grupo investigación UMH 1

Available online: 24 May 2011

To cite this article: J. C. Ospina-E, A. Sierra-C, O. Ochoa, J. A. Pérez-Álvarez & J. Fernández-López (2012): Substitution of Saturated Fat in Processed Meat Products: A Review, Critical Reviews in Food Science and Nutrition, 52:2, 113-122

To link to this article: <http://dx.doi.org/10.1080/10408398.2010.493978>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# Substitution of Saturated Fat in Processed Meat Products: A Review

J. C. OSPINA-E,<sup>1</sup> A. SIERRA-C,<sup>1</sup> O. OCHOA,<sup>1</sup> J. A. PÉREZ-ÁLVAREZ,<sup>2</sup>  
and J. FERNÁNDEZ-LÓPEZ<sup>2</sup>

<sup>1</sup>Industria de Alimentos Zenú S.A.S. Cra 64C N° 104-03. Medellín, Colombia Centro de Investigación y Desarrollo Cárnico CI+D

<sup>2</sup>Universidad Miguel Hernández de Elche. Crta. a Beniel. Km. 3,2. Orihuela, Alicante, España. Grupo REVIV, Generalitat Valenciana y Grupo investigación UMH 1

*The food industry is increasingly directing its efforts to produce and commercialize functional foods where the reduction or even elimination of saturated fat is an important goal. This situation arises from the concern of many institutions and individuals worldwide on the growth of non-transmissible diseases, particularly cardiovascular ones. This article presents a revision of the most important research carried out on processed meat products production and looks at the topic from two principal points of view: the nutritional and technological function of fat and the way in which it is gradually being replaced in the above-mentioned products. Many ingredients have been used to substitute fat but while the results concerning the nutritional composition of the final products are generally acceptable, the sensory aspects are not completely solved. This review emphasizes the use of plastic fats because they allow the highest fat substitution levels during its process and consumption without affecting the product behavior.*

**Keywords** fat, lard, backfat, fat replacer, functional food

## INTRODUCTION

Fat is an undesirable component of meat for most consumers and is responsible for it being regarded as a not very healthy food (Arvanitoyannis and Van Houwelingen-Koukaliaroglou, 2005; Webb and O'Neill, 2008). Health authorities worldwide are encouraging people to decrease their consumption of saturated fatty acids (SFA), *trans* fatty acids, and cholesterol because of many studies which have pointed its direct relation with cardiovascular diseases. At the same time, campaigns encourage people to have balanced diets, take regular physical exercise, and in general assume healthy habits. Such advice has been incorporated into many norms and laws by a variety of organizations whose aim is to promote the health and well-being of people (Pan American Health Organization, 2007; OMS, 2004; FDA/CFSAN, 2004).

One consequence of this is that consumers are more aware of the fact that what they eat is linked directly to their health

(Sampaio et al., 2004). This explains why more consumers are always looking to the nutritional information provided in a product and in many cases this information determines their decision whether to buy it or not.

Also, consumers are showing a preference for functional foods and this must be considered as an important challenge for the meat processing industry. Sausages are rich in SFA and cholesterol; both come from animal fat and have a limited place in a healthy diet. Therefore, there is a need to find fat replacers that will permit the development of meat products with a lower content of SFA, *trans* fatty acids, and cholesterol which at the same time will preserve the sensory qualities and shelf life of the products that guarantee their acceptance by consumers (Sampaio, et al., 2004; Muguerza et al., 2003; 2004).

The aim of this article is to present an overview of the back fat replacement in meat products to fulfil the textural functions that this fat provides to the final product. The fat origin of the products is emphasized because the calorie reduction per se is not the principal objective of the research being carried out. The revision looks at the nutritional and technological function of fat and the way in which it can be substituted in cooked meat products.

Address correspondence to Juana Fernández López, Universidad Miguel Hernández de Elche. Crta. a Beniel. Km. 3,2. Orihuela, Alicante, España. Grupo REVIV, Generalitat Valenciana y Grupo investigación UMH 1 Tel.: 34-966749784 Fax: 34 966749677. E-mail: j.fernandez@umh.es

## NUTRITIONAL AND TECHNOLOGICAL FUNCTION OF FATS

The fats in the diet are composed of triglycerides, glycerol, and fatty acids, which are generally classified according to their chemical structure into saturated (SFA) and unsaturated (UFA), which in turn may be monounsaturated (MUFA) or polyunsaturated (PUFA). These fatty acids, depending on their proportions, define the physical properties and differentiate one fat from the others.

Pig hard fat is the most widely used fat in sausages because of the superior taste and textural characteristic that comes from its higher SFA content. Soft fats, on the other hand, contain a higher proportion of UFA that takes to both technological and sensory problems which persist in the product, where they affect its appearance, make difficult the cutting process, and lead to a greater tendency to oxidation (Maw et al., 2003). These aspects need to be kept in mind when designing or choosing a material of fatty origin to replace those generally present in the meat matrix.

The consumption of pig fat may increase the levels of SFA and cholesterol in the organism, where they are directly related with cardiovascular diseases (Bragagnolo and Rodriguez-Amaya, 2002; FDA/CFSAN, 2004; García-García and Totosaús, 2008; Muguerza et al., 2001; OMS, 2004; Pan American Health Organization, 2007; Williams, 2000; Wood et al., 2003). However, such characteristics are in sharp contrast to their technological qualities which make them indispensable in the manufacture of meat products. The technological and nutritional qualities of backfat are therefore inversely related (Hugo and Roodt, 2007), which means that great care has to be taken when designing a substitute for this material as both criteria must be fulfilled.

### Nutritional Criteria for Good Quality Fats

The fat that is consumed as part of the diet performs various functions in the organism: it provides energy and essential fatty acids, maintains the body temperature constant, protects vital organs, facilitates the absorption of liposoluble vitamins (A, D, E, and K), and promotes the slow evacuation of the stomach, conferring the satiety sensation due to the high energetic density that it provides: 9 cal/g (Pinheiro and Penna, 2004).

Some studies have shown that low fat diets may increase triglyceride levels during fasting and after eating and reduce the protection capacity of High Density Lipoprotein (HDL) cholesterol (Williams, 2000). It is therefore necessary to adjust eating habits to ensure the ingestion of adequate proportions of fatty acids.

The consumption of recommended fat levels in the diet (total fat should be less than 30% and SFA less than 10% of daily total energy intake) will enable the organism to fulfil its functions, while diminishing the risk of cardiovascular diseases (Williams, 2000; Wood et al., 2003). In addition, MUFA have a neutral/

favorable effect on cardiovascular diseases (Warnants et al., 1998; Williams, 2000).

The main parameters studied to determine the nutritional quality of the lipid fraction of the diet are: the SFA/PUFA ratio and *n-3/n-6* PUFA (Pelser et al., 2007; Ansorena and Astiasarán, 2004), as well as the Total Fatty Acids (TFA) and cholesterol content.

### PUFA/SFA Ratio

Fatty acids have different effects according to the concentrations of lipids and lipoproteins in blood plasma. SFA of 12–16 carbons long tend to raise low density lipoprotein (LDL) and total cholesterol levels, the strongest effect has been attributed to lauric (C12:0), myristic (C14:0), and palmitic (C16:0) acids while stearic acid (18:0) has no effect on cholesterol (Risérus et al., 2009; Hu et al., 2001; Valsta et al., 2005; Williams, 2000).

The main health benefits of PUFA intake are associated with the consumption of *n-3* fatty acids. Epidemiological studies and clinical assays have demonstrated that *n-3* PUFAs reduce the incidence of cardiovascular diseases (Kris-Etherton et al., 2002; Wang et al., 2006; Webb and O'Neill, 2008; Williams, 2000).

To improve health it is recommended to eat foods rich in *n-3* PUFA, maintaining the PUFA/SFA ratio greater than 0.4 (Ansorena and Astiasarán, 2004; Hugo and Roodt, 2007; Pelser et al., 2007; Özvural and Vural, 2008; Webb and O'Neill, 2008; Wood et al., 2003). This may be achieved by replacing SFA in the diet by PUFA or MUFA, where PUFA could be more effective in the cholesterol levels reduction but their lipoperoxidation products may have potentially adverse effects (Williams, 2000).

Meat naturally has 0.1 PUFA/SFA ratio which implies an imbalance in the fatty acids intake (Webb and O'Neill, 2008; Wood et al., 2003) while backfat has a value greater than of 0.4 (Gläser et al., 2004; Warnants et al., 1998; Ospina et al., 2010). This means that in the design of a fat replacer attention should be paid to the SFA content in order to improve the PUFA/SFA ratio in the final product.

### PUFA *n-6/n-3* Ratio

The *n-3* and *n-6* fatty acids cannot be synthesized by humans and therefore they are considered essential fatty acids. The *n-6* fatty acids can be elongated as far as arachidonic acid (C20:4 *n-6*) which acts as a precursor in the eicosanoids formation. The role of *n-3* PUFA is important because it inhibits the eicosanoids biosynthesis and reduces inflammation; also, it increases the levels of HDL cholesterol and reduces LDL cholesterol levels (Johnston, 2009; Osborn and Akoh, 2002; Simopoulos, 2000; 2002). This explains why the *n-6/n-3* ratio has been studied as a risk factor in cardiovascular diseases especially via pro-thrombotic and pro-inflammatory pathways that could lead to heart attacks (Fernández-Ginés et al., 2005; Williams, 2000; Wood et al., 2003).

The principal source of *n-6* in the diet is linoleic acid (C18:2) which is found at high concentrations in pork unlike

in other meat species (Wood et al., 2003). The sources of *n*-3 PUFA are  $\alpha$ -linolenic acid (C18:3), eicosapentanoic acid (EPA, C20:5), and docosahexanoic acid (DHA, C22:6). However, it has been demonstrated that the increased consumption of *n*-3 from C18:3 has no beneficial effect on health (Wang et al., 2006).

The *n*-6/*n*-3 western diets ratio is around 15–20 and the recommendation is to eat more foods rich in *n*-3 in order to keep the *n*-6/*n*-3 ratio below 4 (Ansorena and Astiasarán, 2004; Hugo and Roodt, 2007; Pelser et al., 2007; Simopoulos, 2000; Webb and Oá;Neill, 2008; Wood et al., 2003).

This *n*-6/*n*-3 ratio in the meat and fat of animals can be modified by altering their feed if the subsequent meat sensory qualities are not important (Moghadasian, 2008; Ngapo and Gariépy, 2008; Valencia et al., 2007; Wood et al., 2003). However, it is possible to design pig fat substitutes that do not affect the sensory characteristics of the final meat products and improve their *n*-6/*n*-3 ratio.

To summarize, it is not a question of improving the PUFA/SFA ratio by simply adding more PUFA instead of SFA; also, it is important to consider the ratio within this group of *n*-6 and *n*-3 PUFAs.

### *Trans Fatty Acids*

The *trans* fatty acids are some of the isomers that may be formed in the UFA. Despite being unsaturated, *trans* fatty acids behave as SFA due to their spatial conformation, because they form lineal chains around the double bond which may be packed into confined spaces and therefore have similar effects as saturated fatty acids on the organism (Semma, 2002).

Although SFA are mainly responsible for increasing the level of LDL in the diet, *trans* fatty acids and dietetic cholesterol make their own significant contribution (FDA/CFSAN, 2004; Williams, 2000). The direct relation between the consumption of *trans* fatty acids and the risk of cardiovascular diseases may be the result of several mechanisms: increased levels of seric LDL cholesterol and decreased levels of HDL cholesterol, increased plasma levels of triglycerides, inhibition of the  $\Delta$ 6-desaturase enzyme which participates in the metabolism of essential fatty acids, and prostaglandins that may lead to thrombogenesis (Tanasescu et al., 2004; Hu et al., 1999; 2001; Riséru et al., 2009; Williams, 2000).

Besides their negative effects on cardiovascular health, SFA have another harmful effect like inhibiting the essential fatty acids conversion which have important functions in the nervous system development and the correct sight operation since they use the same enzymes to be metabolized (IFST, 2007).

Many laws have attempted to make mandatory mentioning the content of this fatty acid type on labels. It is for this reason that design of any food should consider their inclusion in the formula where they are important not only for their technological function but also for their nutritional contribution.

### *Cholesterol*

Cholesterol is an essential component of animal cell membranes and a precursor of biologically active compounds such as bile salts, important hormones, and vitamin D3 (Morgado, 2008). Although cholesterol is frequently related to risk factors associated with developing cardiovascular diseases, it would be more correct to speak of the association between cholesterol and the HDL and LDL.

The body's cholesterol homeostasis depends on the precise regulation of the cholesterologenesis processes, diet's cholesterol absorption and excretion. Any imbalance through these processes may turn into several health problems like high cholesterol concentrations in plasma, its accumulation in tissues and increased risk of CV diseases (Calpe-Berdiel et al., 2009). In addition, there is a direct relationship between high plasmatic cholesterol levels, especially LDL types, and the risk of cardiovascular diseases (Calpe-Berdiel et al., 2009). LDL reduction seems to stabilize the plaque burden and therefore reduces future cardiovascular events. When the HDL cholesterol is increased, plaque burden could be decreased, intensifying the effectiveness in primary prevention and in advanced atheroma progression (Sirtori and Fumagalli, 2006).

Meat and meat products are important cholesterol sources, they usually provide between 30 and 120 mg of cholesterol per 100 g of food (Valsta et al., 2005). Mixtures of vegetal oils can be designed to ensure the absence of cholesterol in fat replacers and limiting its presence in meat products only to that contributed by the meat itself. This empowers the manufacturer to concentrate on other nutritional fat quality factors when planning a pig fat substitute.

### *Technological Criteria of Good Quality Fat*

Fat is in meat supply animals like subcutaneous deposits as intermuscular and intramuscular forms. Each of these fatty deposits has a particular role in processed meat products. That is why intramuscular fat is particularly desired in the production of ham, intermuscular fat in spreadable products, and subcutaneous fats in the majority of emulsified and mixed meat products.

Backfat is a hard fat used in the manufacturing of meat products and technologically it is preferred to bovine and poultry fats. SFA in pig are present in both meat and lard, the exact proportion varying with the fat present in the feed provided to the animals (Eder et al., 2001; Hugo and Roodt, 2007; Nuernberg et al., 2005; Pascual et al., 2006; Seman, 2008).

Backfat gives textural attributes to meat products such as hardness, gumminess, juiciness, and chewiness, as a result of its physical characteristics that depend on temperature and fatty acid composition (Muguerza et al., 2001; Hsu and Yu, 2002; Fernández-Ginés et al., 2005; Hugo and Roodt, 2007; Wood et al., 2003). Those technological quality attributes are defined by aspects such as: the percentage of extractable fat, firmness, consistency, color, iodine index, C18:2 content, C18:0/C18:2 ratio, the double bond index, fatty acid content, fusion behavior,

solid fat content (SFC), free fatty acid content, oxidative stability, peroxide index, saponification index, crystallization time, among other factors (Wood et al., 2003; Gläser et al., 2004; Hugo and Roodt, 2007; Seaman, 2008).

The aspects that predict the technological attributes of fat, both physical and chemical, always depending on temperature, are shown in Figure 1. It can be seen that three of the most important physical characteristics are crystallization time, melting point, and slip point may lie in the SFC as one of its points, regardless of the method used to measure it. For this reason, the analysis of this variable which determines the quantity of fat in solid and liquid state at a given temperature is more complete. Other chemical indexes are important from the fat lipid oxidation point of view and the consequences of such oxidation in the elaboration of processed meat products.

### Physical Characteristics

Good quality backfat is usually defined as the one which is white and firm, although it is also important to consider its melting behavior.

Chemically, the instauration level of backfat may be sufficient for predicting the firmness behavior of backfats, but not all the aspects are related to the melting point because it also depends on the MUFA and PUFA content. In the same way the consistency of the fat could be predicted by variables such the SFA content particularly C18:0, C18:2, and the ratio between them, the MUFA/SFA ratio and the iodine index (Gläser et al., 2004).

The melting point and pig fat consistency are related to the C18:0 content. Palmytic acid may influence the consistency more at low temperatures when the slip point is used to measure it. When the C18:2 content is greater than 15%, the fat firmness and melting point are determined by this fatty acid and the C16:0 content (Hugo and Roodt, 2007).

Gläser et al. (2004) maintained that although the firmness of backfat depends on the fatty acid make-up only 30% of the variation can be explained by this variable. There are other factors such as the fat, water, and collagen contents, all of which are related and the proportion of surface fat which shows the best

correlation. From these observations, it seems that the quantity of surface fat and its thickness are more important for the firmness than the composition of the backfat.

Consistency might also be related to the SFC. For example, Davenel et al. (1999) suggested that a SFC20 of less than 15% represents a soft fat while a SFC20 of more than 18% represents a hard pig adipose tissue. The SFC is more related with the SFA content than with other fatty acids present in backfat (Davenel et al., 1999; Gläser et al., 2004). Furthermore, the SFA content is directly related with the melting point of fat (Wood et al., 2008). When fat cells have a higher melting point they are in a solid state at room temperature and are more white in color than fats of lower melting point (Wood et al., 2003) which is another characteristic of a good quality fat.

However, color is not the most critical variable when it comes to thinking of pork fat replacers and several studies have pointed out that the use of vegetal oils instead of backfat does not produce significant changes in the color of finished products when assessed by a sensory panel (Muguerza et al., 2001; Pelser et al., 2007).


Lard consistency is then the most interesting physical variable because it is the attribute that will finally gives texture characteristics to the finished product. The relation of other physical variations, especially SFC for predicting how consistency will behave at different temperatures is important when designing a fat replacer; it is also important to consider that the chemical composition will modify the physical lard behavior.

### Chemical Composition

The chemical characteristics evaluated in the lard are the extractable fat percentage and fatty acid profile with particular reference to total C18:2, the C18:0/C18:2 ratio and the double bond index (Hugo and Roodt, 2007). Oxidation is another important quality index associated with the chemical composition since it affects fat color and taste with time. Some of the data studied by other authors and revised by Hugo and Roodt (2007) are shown in Table 1, which includes several indexes based on the chemical composition and associated with the technological criteria defining good quality backfat.

When a comparison is made between the values that define a good nutritional quality backfat ( $\text{PUFA/SFA} > 0.4$  and  $\text{PUFA } n-6/n3 < 4$ ) and the good technological quality indexes (Table 1) it can be seen that it is not possible to maintain the relations suggested from a nutritional point of view ( $\text{PUFA/SFA} = 0.366$  and  $\text{PUFA } n-6/n3 = 4$ ). As more fat is added to the meat product, the nutritional quality decreases and the ratio of the fatty acids gets worse.

Considering the large number of criteria necessary to define lard quality, in order to get a good backfat substitute, the most representative physical characteristics should be preserved while the chemical characteristics should be modified to reach the nutritional criteria identified as good quality fat indexes.



**Figure 1** Physical and chemical parameters used as quality criteria in backfat. (color figure available online.)

## SUBSTITUTING FAT IN MEAT PRODUCTS

Meat products are ripe candidates for being modified and turned in healthy products through the addition of ingredients considered beneficial for health or through the elimination or reduction of those components considered harmful (Fernández-Ginés et al., 2005). Among these components fat must be included as one of the factors that could be used to adjust the nutritional characteristics of the product.

The substitution of fat has been studied for many years and has its own terminology. For example, Akoh (1998) coined the term “fat replacers” to denote the ingredients or additives used to diminish the fat content in foods. These are ingredients which structurally may be considered fats, proteins, or carbohydrates, that is, any raw material used to replace fat in the formulation. Within the group of “fat replacers” there are two sub-groups: “fat mimetics” and “fat substitutes.”

### Fat Mimetics

These are substances that mimic the physical and organoleptic triglycerides characteristics but cannot replace the fat on a 1:1 basis. Basically these types of ingredients are carbohydrates and proteins that require hydration to fulfil their function as fat replacers. However, they have their drawbacks: for example, they undergo caramelization and denaturalization when they are submitted to high temperatures; they have soluble but not lipidic flavors so the taste is less and generally require additional preparation processes. Their main advantage is that Fat Mimetics reduce food caloric content (Akoh, 1998).

Partial substitution of backfat with Fat Mimetics has been developed in order to reduce the negative effect of the high fat content in meat products. In several studies where fat has been partially replaced by fat mimetics the ingredients used were water, proteins, and polysaccharides like maltodextrins,

gums, carragenins, fibers, inulin, starch, flours, bran, and even tubers (Andrès et al., 2006; Cengiz and Gokoglu, 2007; Cierach et al., 2009; Crehan et al., 2000; García et al., 2002, 2006; García-García and Totosa, 2008; Fernández-Ginés et al., 2005; Mendoza et al., 2001; Nowak et al., 2007; Piñero et al., 2008; Salazar et al., 2009; Sampaio et al., 2004; Shon and Chin, 2008; Tan et al., 2007).

### Fat Substitutes

These are macromolecules that physically and chemically resemble triglycerides and so they can be expected to be used as fat substitutes in a 1:1 proportion. Generally they are elaborated from conventional fatty acid bases that are physically and chemically altered (Akoh, 1998). Animal fat substitutes made from vegetal oils have gained much attention in the processed meat industry. Such fat substitutes can be classified into two groups: liquids and plastics (Tan et al., 2006).

Liquid fats are basically represented by vegetal oils and have been seen to have a positive impact on nutritional aspects as represented by reduced cholesterol content and improved PUFA/SFA and *n-6/n-3* ratios. Sunflower, maize, peanut, tea seed, coconut, palm, soy, and olive oils as well as fish oil have been evaluated (Hsu and Yu, 2002; Fernández-Ginés et al., 2005; López-López et al., 2009; Moghadasian, 2008; Muguerza et al., 2001; 2002; 2003; 2004; Severini et al., 2003; Yildiz-Turp and Serdaroglu, 2008).

Having in mind the proportions mentioned above, the use of vegetal oils can be compared with the use of soft fats which are characterized by their poor appearance, difficulty to cut, and greater tendency to oxidize than hard fats (Maw et al., 2003). This tendency to oxidation in the products containing vegetal oils instead of lard is due to the greater unsaturated fatty acid content, that is why it is vital to define the shelf life that results from increased rancidity and decoloration (Fernández-Ginés et al., 2005; Hugo and Roodt, 2007; Webb and O'Neill, 2008).

Plastic fats are obtained by chemically and enzymatically altering some oils; they appear to be solid and are resistant to light forces but when these forces pass a certain value plastic fats flow like liquids (Osborn and Akoh, 2002). Partial hydrogenation and interesterification are used to modify oils in an attempt to simulate the consistency and melting point of animal fat. Interesterification is a healthy alternative because it does not cause saturation of the fatty acids and does not form *trans* fatty acids (Özvural and Vural, 2008; Vural et al., 2004). Animal fats have been replaced with a variety of vegetal oils including palm, cotton, olive, hazelnut, and their mixtures (Özvural and Vural, 2008; Vural et al., 2004).

The most recent studies into the replacement of animal fats in meat products are depicted in Table 2. The studies involving plastic fats are remarkable instead of the traditional use of vegetal oils and their pre-emulsified forms.

**Table 1** Technological criteria defining a good quality lard (adapted from Hugo and Roodt, 2007)

Variable	% min	% max
Composition		
Extractable fat	84	90
SFA	41	—
UFA	—	59
MUFA	—	57
PUFA	—	15
C18:0	12	—
C18:2	12	15
C18:0/C18:2	1.2	—
Oxidation		
PUFA	—	23
dienoic fatty acid	—	10
trienoic fatty acid	—	1.0
tetraenoic fatty acid	—	0.5
pentaenoic +hexaenoic fatty acid	—	1.0
Double bonds index	—	80

Table 2 Fat substitution in processed meat products

Authors	Product	Add Fat (%)	Substitution (%)	Substitute	Sensory Prop. (acceptability)					Color			TPA					Author Conclusion	
					Appearance	Texture	Taste	Odour	Color	Overall	L*	a*	b*	Hardness	Cohesiveness	Gumminess	Springiness		Chewiness
Choi et al. (2010)	Frankfurters	30	50*	OO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes
			50*	GSO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	CO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	KO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	SO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Choi et al. (2009)	Meat batter emulsified	30	50*	OO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	GSO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	CO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	KO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50*	SO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Martín et al. (2008)	Pork Liver Paté		50*	OO + SC (PE) + RBF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50	OO + SC (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			30	CLA + SC (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
Özvural and Vural (2008)	Frankfurters	30	50	OO50% + CLA50% + SC (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			10	PO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	PS	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Yildiz-Turp and Serdaroğlu (2008)	Turkish fermented sausage (Sucuk)		100	CSO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	HO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	33.3%PO + 33.3%PS + 33.3%HO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	33.3%PO + 33.3%PS + 33.3%CSO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	16.6%PO + 16.6%PS + 66.6%HO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Yildiz-Turp and Serdaroğlu (2008)	Turkish fermented sausage (Sucuk)		100	16.6%PO + 16.6%PS + 66.6%CSO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	8.3%PO + 8.3%PS + 83.4%HO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			100	8.3%PO + 8.3%PS + 83.4%CSO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			15	HO + WP (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			20		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Valencia et al. (2007)	Chorizo Pamplona	25	30		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			50	Algae Schizochytrium sp. Oil	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			25		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
Pelser et al. (2007)	Dutch style fermented sausage	30	10	FSO + SP (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			15	FSO + SP (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			20	FSO + SP (PE)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
Tan et al. (2006)	Chicken Frankfurters		20	FSO + SP (SC)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			15	Encapsulated FSO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			15	Encapsulated FO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			50	PS	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			100	PS	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
Tan et al. (2006)	Chicken Frankfurters		100	PS50% + PO50%	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			100	PO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			50	PO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			66.6	PS 50% + PO50%	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			33.3	PS 50% + PO50%	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
Tan et al. (2006)	Chicken Frankfurters		83	PS80% + PO20%	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	No	
			83	PS20% + PO80%	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	
			83		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Yes	

(Continued on next page)

Table 2 Fat substitution in processed meat products (Continued)

Authors	Product	Add Fat (%)	Substitution (%)	Substitute	Sensory Prop. (acceptability)								Color			TPA					Author Conclusion									
					Cooking loss	Consistency	Appearance	Texture	Taste	Odour	Color	Overall	L*	a*	b*	Hardness	Cohesiveness	Gumminess	Springiness	Chewiness		TBA Value								
Valencia et al. (2006) Ansorena and Astiasarán (2004)	Chorizo Pamplona	25	100	Deodorized FO + SP (PE)																										Yes
	Chorizo Pamplona	25	25	LO + SP (PE)																										Yes
Vural et al. (2004)	Frankfurters	10	25	LO + SP + BHA + BHT (PE)																										Yes
		60	100	Intersterified PO + SBF																										Yes
		60	100	Intersterified PO + SBF																										Yes
		60	100	Intersterified CSO + SBF																										Yes
		60	100	Intersterified CSO + SBF																										Yes
		60	100	Intersterified OO + SBF																										Yes
Severini et al. (2003)	Salami	15	33.3	OO + SC (PE)																										Yes
		50	66.6																											No
Muguerza et al. (2003)	Chorizo Pamplona	25	15	SO + SP (PE)																										Yes
		20	25																											Yes
Hsu & Yu (2002)	Low Fat Kung-Wans	25	0**	BF																										N/A
		25	100**	W																										N/A
		25	100**	CNO																										Yes
		25	100**	PO																										Yes
		25	100**	SO																										Yes
		25	100**	SFO																										No
		25	100**	CO																										No
		25	100**	PNO																										No
		25	100**	TO																										No
		25	100**	OO																										Yes
		25	100**	Hydrogenated CNO																										No
Muguerza et al. (2001)	Chorizo Pamplona	25	100**	Hydrogenated PO																										Yes
		25	100**	Hydrogenated SO																										Yes
		25	10	OO + SP (PE)																										Yes
		15																												Yes
		20																												Yes
			25																										Yes	
			30																										No	

\* The total fat content is also reduced to 20%

\*\* The total fat content is also reduced to 10%

≈ without statistical significance (p > 0.05), except Severini et al. (2008) with p > 0.08; < less than control, > higher than control

OO: Olive Oil, GSO: Grape Seed Oil, CO: Corn Oil, KO: Canola Oil, SO: Soybean Oil, PO: Palm Oil, PS: Palm Stearin, CSO: Cottonseed Oil, HO: Hazelnut Oil, FSO: Flaxseed Oil, FO: Fish Oil, BF: Back Fat, W: Water, CNO: Coconut Oil, SFO: Sunflower Oil, PNO: Peanut Oil, TO: Tea Seed Oil, LO: Linseed Oil, CLA: Conjugated Linoleic Acid, PE: Pre-emulsified, SC: Sodium Caseinate, SP: Soybean Protein, WP: Whey Protein, RBF: Rice Bran Fiber, SBF: Sugarbeet Fiber



Studies where pig fat was substituted by interesterified vegetal oils, 100% substitution has been reached in products where the fat content does not exceed 10% of the formula with good sensory results (Özvural and Vural, 2008; Vural et al., 2004). However, some products have a greater percentage of fat which means that solutions have to be found to the problem this represents.

Another alternative included in Table 2 is the use of vegetal or animal oils as individual substitute in products with an added fat content lower than 10%, for which good organoleptic results have been obtained (Hsu and Yu, 2002; Özvural and Vural, 2008; Pelsner et al. 2007; Tan et al. 2006; Valencia et al., 2007). However, as the level of fat increases in the formula so the level of substitution falls; for example, substitutions of more than 25% of the fat in products containing more than 20% added fat have not been successful.

The use of intermediate processes such as pre-emulsions using different types of protein represents an improved way of incorporating oil in meat matrices, with this higher levels of fat substitution have been obtained than those with the direct use of oils (Ansorena and Astiasarán, 2004; Choi et al., 2010; 2009; Martin et al., 2008; Muguerza et al., 2001; 2003; Pelsner et al., 2007; Severini et al., 2003; Valencia et al., 2006; Yildiz-Turp and Serdaroğlu, 2008). However, no substitutions above 25% have been possible when fat exceeds 25%. Moreover, from an industrial point of view this type of solution is not interesting because it would affect productivity especially in cooked products cases.

The results obtained in measuring different response variables (Table 2) indicate that it is the sensory characteristics that finally define the fat substitution success, although instrumental color measures and textural and lipid oxidation analysis are also objective evaluations to get closer to the organoleptic profile and provides a more complete evaluation to compare the results from different studies (Ngapo and Gariépy, 2008), even though it is necessary to establish a hierarchy or an evaluations system to conclude.

## CONCLUSIONS

By following the nutritional indexes suggested throughout this revision, meat products can be considered as neutral character food that not reduces or increases the risk of cardiovascular diseases. However, fat is not the only source of fatty acids in processed meat products and meat itself may be a contributor to the deterioration of the suggested indices. Moreover, meat products are not the only components of the human diet and other sources of fat should be borne in mind to maintain the balance suggested. Finally, the risk of cardiovascular diseases is associated with eating habits in general, lifestyle, and even has a genetic component. The meat industry's responsibility should be at least to offer products that do not increase the risk of appearance of non-transmissible diseases.

Loads of fat replacer have been evaluated to reduce the fat added to meat products; however, the ideal one has not yet been identified. The use of plastic fats as backfat substitutes permits high levels of substitution, sensory attributes to be maintained, and the nutritional characteristics of processed meat products to be improved. The use of mixtures of chemically modified vegetal oils should be explored for products with a high level of added fat.

## REFERENCES

- Andrès, S., Zaritzky, N., and Califano, A. (2006). The effect of whey protein concentrates and hydrocolloids on the texture and color characteristics of chicken sausages. *International Journal of Food Science and Technology*. **41**: 954–961.
- Ansorena, D. and Astiasarán, I. (2004). The use of linseed oil improves nutritional quality of the lipid fraction of dry-fermented sausages. *Food Chemistry*. **87**: 96–74.
- Akoh, C.C. (1998). Fat replacers. *Food Technology*. **52**:47–53.
- Arvanitoyannis, I. S. and Van Houwelingen-Koukarioglou, M. (2005). Functional foods: A survey of health claims, pros and cons, and current legislation. *Critical Reviews in Food Science and Nutrition*. **45**: 385–404.
- Bloukas, J.G., Paneras E.D., and Fournitzis, G.C. (1997). Effect of replacing pork backfat with olive oil on processing and quality characteristics of fermented sausages. *Meat Science*, **45**: 133–144.
- Bragagnolo, N. and Rodriguez-Amaya D. (2002). Simultaneous determination of total lipid, cholesterol and fatty acids in meat and backfat of suckling and adult pigs. *Food Chemistry*. **79**: 255–260.
- Calpe-Berdiel, L., Escolà-Gil, J.C., and Blanco-Vaca, F. (2009). Review: New insights into the molecular actions of plant sterols and stanols in cholesterol metabolism. *Atherosclerosis*. **203**: 18–31.
- Cengiz, E. and Gokoglu, N. (2007). Effects of fat reduction and fat replacer addition on some quality characteristics of frankfurter-type sausages. *International Journal of Food Science and Technology*. **42**: 366–372.
- Choi, Y.S., Choi, J.H., Han, D.J., Kim, H.Y., Lee, M.A., Jeong, J.Y., Chung, H.J., and Kim, C.J. (2010). Effects of replacing pork back fat with vegetable oils and rice bran fiber on the quality of reduced-fat frankfurters. *Meat Science*. **84**: 557–563.
- Choi, Y.S., Choi, J.H., Han, D.J., Kim, H.Y., Lee, M.A., Kim, H.W., Jeong, J.Y., and Kim, C.J. (2009). Characteristics of low-fat meat emulsion systems with pork fat replaced by vegetable oils and rice bran fiber. *Meat Science*. **82**: 266–271.
- Crehan, C.M., Hughes, E., Troy, D.J., and Buckley, D.J. (2000). Effects of fat level and maltodextrin on the functional properties of frankfurters formulated with 5, 12, and 30% fat. *Meat Science*. **55**: 463–469.
- Cierach, M., Modzelewska-Kapitula, M., and Szacilo, K. (2009). The influence of carrageenan on the properties of low-fat frankfurters. *Meat Science*. **82**: 295–299.
- Davenel, A., Riaublanc, A., Marchal, P., and Gandemer G. (1999). Quality of pig adipose tissue: relationship between solid fat content and lipid composition. *Meat Science*. **51**: 73–79.
- Eder, K., Nonn, H., and Kluge, H. (2001). The fatty acid composition of lipids from muscle and adipose tissues of pigs fed various oil mixtures differing in their ratio between oleic acid and linoleic acid. *Eur. J. Lipid Sci. Technol*. **103**: 668–676.
- FDA/CFSAN Office of Nutritional Products, Labeling and Dietary Supplements. (2004). Los ácidos grasos trans ahora serán listados junto con las grasas saturadas y el colesterol en las etiquetas de información nutricional. <http://www.cfsan.fda.gov/~dms/stransfa.html#whatis>
- Fernández-Ginés, J.M., Fernández-López, J., Sayas-Barberá, E. and Pérez-Álvarez, J.A. (2005). Meat products as functional foods: A review. *Journal of Food Science*, **70**: R37–R43.

- García, M. L., Cáceres E. and Selgas, M. D. (2006). Effect of inulin on the textural and sensory properties of mortadella, a Spanish cooked meat product. *International Journal Food Science Technology*. **41**: 1207–1215.
- García, M.L., Domínguez, R., Galvez, M.D., Casas, C., and Selgas M.D. (2002). Utilization of cereal and fruit fibers in low fat dry fermented sausages. *Meat Science*, **60**: 227–236.
- García-García, E. and Totosa, A. (2008). Low-fat sodium-reduced sausages: Effect of the interaction between locust bean gum, potato starch and *k*-carrageenan by a mixture design approach. *Meat Science*, **78**: 406–413.
- Gläser, K., Wenk, C., and Scheeder, M. (2004). Evaluation of pork backfat firmness and lard consistency using several different physicochemical methods. *Journal of the Science of Food and Agriculture*. **84**: 853–852.
- Hsu, S.Y and Yu, S.H. (2002). Comparisons on 11 plant oil fat substitutes for low-fat Kung-wans. *Journal of Food Engineering*. **51**: 215–220.
- Hu, F.B., Manson, J.E.Y., and Willett, W.C. (2001). Types of dietary fat and risk of coronary heart disease: A critical review. *Journal of the American College of Nutrition*, **20**: 5–19.
- Hu, F.B., Stampfer, M.J., Manson, J.E., Ascherio, A., Colditz, G.A., Speizer, F.E., Hennekens, C.H.Y., and Willett, W.C. (1999). Dietary saturated fats and their food sources in relation to the risk of coronary heart disease in women. *The American Journal of Clinical Nutrition*. **70**: 1001–1008.
- Hugo, A. and Roodt, E. (2007). Significance of porcine fat quality in meat technology: A review. *Food Review International*, **23**: 175–198.
- Institute of Food Science and Technology IFST. (2007). Trans Fatty Acids (TFA) Information Statement.
- Johnston, C. (2009). Functional foods as modifiers of cardiovascular disease. *American Journal of Lifestyle Medicine*, **3**: 39s–43s.
- Kris-Etherton, P.M., Harris, W.S., Appel, L.J., and for the Nutrition Committee Circulation (2002). Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*. **106**: 2747–2757.
- López-López, I., Cofrades, S., and Jiménez-Colmenero, F. (2009). Low-fat frankfurters enriched with n-3 PUFA and edible seaweed: Effects of olive oil and chilled storage on physicochemical, sensory and microbial characteristics. *Meat Science*. **83**: 148–154.
- Martin, D., Ruiz, J., Kivikari, R., and Puolanne, E. (2008). Partial replacement of pork fat by conjugated linoleic acid and/or olive oil in liver pâtés: Effect on physicochemical characteristics and oxidative stability. *Meat Science*. **80**: 496–504.
- Maw, S.J., Fowler, V.R., Hamilton, M., and Petchey, A.M. (2003). Physical characteristics of pig fat and their relation to fatty acid composition. *Meat Science*. **63**: 185–190.
- Mendoza, E., García, M.L., Casas, C., and Selgas, M.D. (2001). Inulin as fat substitute in low fat, dry fermented sausages. *Meat Science*. **57**: 387–393.
- Moghadasian, M. H. (2008). Advances in dietary enrichment with N-3 fatty acids. *Critical Reviews in Food Science and Nutrition*. **48**: 402–410.
- Morgado, N. (2008). El colesterol: Digestión, absorción, transporte, y Metabolismo extra e intracelular. Diplomado en Grasas y Aceites en la Nutrición Humana. Instituto de Nutrición y Tecnología de los Alimentos (INTA), Versión 3. **8**: 1–34.
- Muguerza, E., Ansorena, D., and Astiasarán, I. (2003). Improvement of nutritional properties of Chorizo de Pamplona by replacement of pork backfat with soy oil. *Meat Science*. **65**: 1361–1367.
- Muguerza, E., Fista, G., Ansorena, D., Astiasarán, I., and Bloukas, J.G. (2002). Effect of fat level and partial replacement of pork backfat with olive oil on processing and quality characteristics of fermented sausages. *Meat Science*. **61**: 397–404.
- Muguerza, E., Gimeno, O., Ansorena, D., and Astiasarán, I. (2004). New Formulations for healthier dry fermented sausages: A review. *Food Science and Technology*. **15**: 452–457.
- Muguerza, E., Gimeno, O., Ansorena, D., Bloukas, J.G. and Astiasarán, I. (2001). Effect of replacing pork backfat with pre-emulsified olive oil on lipid fraction and sensory quality of Chorizo de Pamplona: A traditional Spanish fermented sausage. *Meat Science*. **59**: 251–258.
- Ngapo, T. M. and Gariépy, C. (2008). Factors affecting the eating quality of pork. *Critical Reviews in Food Science and Nutrition*. **48**: 599–633.
- Nowak, B., Von Mueffling, T., Grotheer, J., Klein, G., and Watkinson, B.-M. (2007). Energy content, sensory properties, and microbiological shelf life of German bologna-type sausages produced with citrate or phosphate and with inulin as fat replacer. *Journal of Food Science*. **72**: S629–S638.
- Nuernberg, K., Fischer, K., Nuernberg, G., Kuechenmeister, U., Klosowska, D., Eliminowska-Wenda, G., Fiedler, I., and Ender, K. (2005). Effects of dietary olive and linseed oil on lipid composition, meat quality, sensory characteristics and muscle structure in pigs. *Meat Science*. **70**: 63–74.
- Organización Mundial de la Salud, OMS. (2004). Estrategia Mundial Sobre el Régimen Alimentario, Actividad Física y Salud.
- Osborn, H.T. and Akoh, C.C. (2002). Structured lipids: Novel fats with medical, nutraceutical and food applications. *Comprehensive Reviews in Food Science and Food Safety*. **1**: 110–120.
- Ospina-E, J.C., Cruz-S, A., Pérez-Álvarez, J.A., and Fernández-López, J. (2010). Development of combinations of chemically modified vegetable oils as pork backfat substitutes in sausages formulation. *Meat Science*. **84**: 491–497.
- Özvural, E.B. and Vural, H. (2008). Utilization of interesterified oil blends in the production of frankfurters. *Meat Science*. **78**: 211–216.
- Pan American Health Organization. (2007). Las Américas libres de grasas Trans.
- Pascual, J.V., Rafecas, M., Canela, M.A., Boatella, J., Bou, R., Barroeta, A.C., and Codony, R. (2006). Effect of increasing amounts of a linoleic-rich dietary fat on the fat composition of four pig breeds. Part II: Fatty acid composition in muscle and fat tissues. *Food Chemistry*. **100**: 1639–1648.
- Pelser, M., Linssen, J., Legger, A., and Houben, J. (2007). Lipid oxidation in n-3 fatty acid enriched Dutch style fermented sausages. *Meat Science*. **75**: 1–11.
- Pinheiro M. and Penna A. (2004). Substitutos de gordura: Tipos e aplicações em produtos lácteos. *Alimentos e Nutrição Araraquara*. **15**: 175–186.
- Piñero, M. P., Parra, K., Huerta-Leidenz, N., Arenas de Moreno, L., Ferrer, M., Araujo, S., and Barboza, Y. (2008). Effect of oat's soluble fiber ( $\beta$ -glucan) as a fat replacer on physical, chemical, microbiological and sensory properties of low-fat beef patties. *Meat Science*. **80**: 675–680.
- Risérus, U., Willett, W.C., and Hu, F.B. (2009). Dietary fats and prevention of type 2 diabetes. *Progress in Lipid Research*. **48**: 44–51.
- Salazar, P., García, M.L., and Selgas, M.D. (2009). Short-chain fructooligosaccharides as potential functional ingredient in dry fermented sausages with different fat levels. *International Journal of Food Science and Technology*. **44**: 1100–1107.
- Sampaio, G., Castellucci, C., Pinto e Silva, M., and Torres, E. (2004). Report: Effect of fat replacers on the nutritive value and acceptability of beef frankfurters. *Journal of Food Composition and Analysis*. **17**: 469–474.
- Selgas, M. D., Cáceres E., and García, M. L. (2005). Long-chain soluble dietary fiber as functional ingredient in cooked meat sausages. *Food Science Technology International*. **11**: 41–47.
- Seman, D.L. (2008). Pork fat quality: A processor's perspective. *Proceedings of the Reciprocal Meat Conference*. **61**: 1–7.
- Semma, M. (2002). Trans fatty acids: Properties, benefits and risks. *Journal of Health Science*, **48**: 7–13.
- Severini, C., de Pilli, T., and Baiano, A. (2003). Partial substitution of pork backfat with extra-virgin olive oil in “salami” products: Effects on chemical, physical and sensorial quality. *Meat Science*, **64**: 323–331.
- Shon, J. and Chin, K.B. (2008). Effect of whey protein coating on quality attributes of low-fat, aerobically packaged sausage during refrigerated storage. *Journal of Food Science*. **73**: C469–C475.
- Simopoulos, A.P. (2000). Human requirement for N-3 polyunsaturated fatty acids. *Poultry Science*, **79**: 961–970.
- Simopoulos, A.P. (2002). Dossier: Polyunsaturated fatty acids in biology and diseases. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomedicine and Pharmacotherapy*. **56**: 365–379.
- Sirtori, C.R. and Fumagalli, R. (2006). LDL-cholesterol lowering or HDL-cholesterol raising for cardiovascular prevention: A lesson from cholesterol turnover studies and others. *Atherosclerosis*, **186**: 1–11.
- Tan, S.S., Aminah, A., Zhang, X.G., and Abdul, S.B. (2006). Optimizing palm oil and palm stearin utilization for sensory and textural properties of chicken frankfurters. *Meat Science*. **72**: 387–397.

- Tan, F.J., Liao, F.Y., Jhan, Y.J., and Liu, D.C. (2007). Effect of replacing pork backfat with yams (*Dioscorea alata*) on quality characteristics of Chinese sausage. *Journal of Food Engineering*. **79**: 858–863.
- Tanasescu, M., Cho, E., Manson, J.E., and Hu, F.B. (2004). Dietary fat and cholesterol and the risk of cardiovascular disease among women with type 2 diabetes. *The American Journal of Clinical Nutrition*. **79**: 999–1005.
- Sirtori, C.R. and Fumagalli, R. (2006). Review: LDL-cholesterol lowering or HDL-cholesterol raising for cardiovascular prevention. A lesson from cholesterol turnover studies and others. *Atherosclerosis*. **186**: 1–11.
- Valencia, I., Ansorena, D., and Astiasarán, I. (2006). Nutritional and sensory properties of dry fermented sausages enriched with *n*-3 PUFAs. *Meat Science*. **72**: 727–733.
- Valencia, I., Ansorena, D., and Astiasarán, I. (2007). Development of dry fermented sausages rich in docosahexaenoic acid with oil from the microalgae *Schizochytrium* sp.: Influence on nutritional properties, sensorial quality and oxidation stability. *Food Chemistry*. **104**: 1087–1096.
- Valsta, L.M., Tapanainen, H., and Männistö, S. (2005). Review: Meat fats in nutrition. *Meat Science*. **70**: 525–530.
- Vural, H., Javidipour, I., and Ozbas, O.O. (2004). Effects of interesterified vegetable oils and sugarbeet fiber on the quality of frankfurters. *Meat Science*. **67**: 65–72.
- Wang, C., Harris, W.S., Chung, M., Lichtenstein, A.H., Balk, E.M., Kupelnick, B., Jordan, H.S., and Lau, J. (2006). *n*-3 fatty acids from fish or fish-oil supplements, but not  $\alpha$ -linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary- prevention studies: A systematic review. *The American Journal of Clinical Nutrition*. **84**: 5–17.
- Warnants, N., Van Oeckel, M. J., and Boucqui, CH. V. (1998). Effect of incorporation of dietary polyunsaturated fatty acids in pork backfat on the quality of salami. *Meat Science*. **49**: 435–445.
- Webb, E.C. and O'Neill, H.A. (2008). The animal fat paradox and meat quality. *Review. Meat Science*. **80**: 28–36.
- Williams, C.M. (2000). Dietary fatty acids and human health. *Annales Zootechnie*. **49**: 165–180.
- Wood, J.D., Richardson, R.I., Nute, G.R., Fisher, A.V., Campo, M.M., Kasapidou, E., Sheard, P.R., and Enser, M. (2003). Effects of fatty acids on meat quality: A review. *Meat Science*. **66**: 21–32.
- Wood, J.D., Richardson, R.I., Nute, G.R., Whittington, F.M., Hughes, S.I., and Hallett, K.G. (2008). Factors controlling fatty acid composition and meat quality in pork and other meats. *Proceedings of the Reciprocal Meat Conference*. **61**: 1–7.
- Yildiz-Turp, G. and Serdaroglu, M. (2008). Effect of replacing beef fat with hazelnut oil on quality characteristics of sucuk: A Turkish fermented sausage. *Meat Science*. **78**: 447–454.