

Beet Sugar Syrup and Molasses as Low-Cost Feedstock for the Enzymatic Production of Fructo-oligosaccharides

IRAJ GHAZI, LUCIA FERNANDEZ-ARROJO, ARANZAZU GOMEZ DE SEGURA,[†]
MIGUEL ALCALDE, FRANCISCO J. PLOU,* AND ANTONIO BALLESTEROS

Departamento de Biotálisis, Instituto de Catálisis y Petroleoquímica, CSIC, Cantoblanco,
28049 Madrid, Spain

Sugar syrup and molasses from beet processing containing 620 and 570 mg/mL sucrose, respectively, were assayed as low-cost and available substrates for the enzymatic synthesis of fructo-oligosaccharides (FOSs). A commercial pectinase (Pectinex Ultra SP-L, from *Aspergillus aculeatus*) characterized by the presence of a transfructosylating activity was used as a biocatalyst. The FOS production increased when lowering the initial pH value of syrup (7.5) and molasses (8.9) to 5.5. Sugar syrup and molasses were diluted in order to reduce substrate viscosity; interestingly, the percentage of FOS with regards to total sugars remained almost constant, which indicated a high transferase-to-hydrolase ratio for this enzyme. Kinetics of FOS production was analyzed. Using approximately 10 U transfructosylating activity per g sucrose, the FOS concentration reached a maximum of 388 mg/mL after 30 h using syrup and 235 mg/mL in 65 h with molasses. These values corresponded to approximately 56 and 49% (w/w), respectively, of the total amount of carbohydrates in the mixture. The enzyme was also covalently immobilized on an epoxy-activated polymethacrylate-based polymer (Sepabeads EC-EP5). We found that immobilized Pectinex Ultra SP-L can be efficiently applied to the synthesis of FOS using syrup and molasses as substrates.

KEYWORDS: β -Fructofuranosidase; fructosyltransferase; transglycosidase; sucrose; prebiotics; poly-methacrylate; Pectinex Ultra SP-L; animal feed

INTRODUCTION

Fructo-oligosaccharides (FOSs) of the inulin type constitute one of the most established groups of prebiotics in the world (1). The addition of such molecules to food products may help to prevent illness, control calcium balance, and contribute to the reduction of antibiotic consumption (2, 3).

FOSs of the inulin type are fructose oligomers with a terminal glucose group, in which 2–4 fructosyl moieties are linked via $\beta(1\rightarrow2)$ -glycosidic bonds (4). Commercial FOSs are mainly composed of 1-kestose (GF₂), nystose (GF₃), and 1^F-fructosyl-nystose (GF₄). The properties of other types of FOSs, such as those of the levan type (5) and the neo-FOSs (6), are very promising, but they are not yet commercially available.

FOSs are industrially produced through fructosyl transfer from pure sucrose using a fungal enzyme (7). FOS-synthesising enzymes are present in many higher plants (asparagus, chicory, onion, Jerusalem artichoke, etc.) and microorganisms, especially fungi (*Aureobasidium pullulans*, *Aspergillus niger*, *Aspergillus oryzae*, etc.) (4, 8–10). FOS-producing enzymes are assigned as hydrolases (β -fructofuranosidases, EC 3.2.1.26) or fructosyl-

transferases (transfructosidases, EC 2.4.1.9). The maximal FOS production for a particular enzyme depends on the relative rate of transfructosylation and hydrolysis (11).

In addition to their use in human food products, the administration of FOS to the diets of some animals results in improvement in feed efficiency and a reduction of diarrhoea and smell in feces (12). To extend the use of FOSs as animal feed additives, it is necessary to minimize production costs. Beet sugar syrup and molasses are cheap and available sources of sucrose and are adequate feedstock for FOS production of animal-feed grade.

Pectinex Ultra SP-L is a commercial enzyme preparation from *Aspergillus aculeatus* used in the food industry for fruit juice processing to reduce viscosity. It contains different pectinolytic and cellulolytic enzymes (e.g., endo-poly-galacturonase, endo-pectinylase, and pectin esterase) (13). Interestingly, Pectinex Ultra SP-L also contains a transfructosylating activity (14, 15). In this work, we have assayed Pectinex Ultra SP-L, soluble or immobilized on a epoxy-activated polymethacrylate (Sepabeads EC-EP), for transformation of beet syrup and molasses into FOSs.

EXPERIMENTAL PROCEDURES

Materials. The beet sugar syrup and molasses were kindly donated by Azucarera Ebro (Valladolid, Spain). Pectinex Ultra SP-L (batch No.

* To whom correspondence should be addressed. Fax: +34-91-5854760. E-mail: fplou@icp.csic.es.

[†] Present address: Facultad de Veterinaria, Universidad Autónoma de Barcelona, Departamento de Ciencia Animal y de los Alimentos, Unidad de Nutrición Animal, 08193 Bellaterra, Barcelona, Spain.

Table 1. Some Specifications of the Beet Sugar Syrup and Molasses Used in This Work

	sugar syrup	molasses
Brix units ^a	69.3	75.1
pH	7.5	8.9
density (g/mL)	1.30	1.40
sucrose (%) ^b	47.5	40.7
[sucrose] (mg/mL) ^c	620	570
betaine (%) ^d	0.6	5.6

^a Grams of dry matter per 100 g of sample, provided by the manufacturer.

^b Grams of sucrose per 100 g of sample, determined by HPLC. ^c Determined by HPLC. ^d Grams of betaine per 100 g of sample, provided by the manufacturer.

KRN05401) was kindly donated by Novozymes A/S. Sepabeads EC-EP5 (batch no. E407P094) were kindly provided by Resindion S.R.L. (Mitsubishi Chemical Corporation, Milan, Italy). Glucose and dinitrosalicylic acid (DNS) were from Sigma. Sucrose and fructose were purchased from Merck. 1-Kestose and nystose were from TCI Europe (Zwijndrecht, Belgium). 1^F-Fructosyl-nystose was from Megazyme (County Wicklow, Ireland). All other reagents and solvents were of the highest available purity and used as purchased.

Standard Activity Microassay. The enzymatic activity toward sucrose was measured following the initial rate of reducing sugars production by the DNS method. The spectrophotometric assay was adapted to a 96 well microplate scale as described (16). One unit (U) of activity was defined as that catalyzing the formation of 1 μ mol reducing sugar per minute under the above conditions.

Immobilization of Fructosyltransferase. Pectinex Ultra SP-L (20 mL) and Sepabeads EC-EP5 (8 g) were mixed and incubated for 24 h at room temperature with roller shaking. The ratio protein/support was approximately 45 mg protein per gram of carrier. The biocatalyst was then filtered, washed (3 \times 30 mL) with 50 mM sodium acetate buffer (pH 5.6), dried under vacuum, and stored at 4 $^{\circ}$ C.

Batch Production of FOSs. Soluble or immobilized Pectinex Ultra SP-L was added to diluted beet sugar syrup and molasses, previously filtered through a glass microfiber filter. The total reaction volume was 5 mL. The biocatalyst was added to a final activity in the mixture of 1 or 5 U/mL (determined by the standard DNS microassay). The mixtures were incubated at 60 $^{\circ}$ C in an orbital shaker (Stuart Scientific) at 200 rpm. At different times, 40 μ L aliquots were extracted from the reaction mixture, diluted with 160 μ L of water, and incubated for 10 min at 90 $^{\circ}$ C to inactivate the enzyme. Samples were centrifuged for 5 min at 6000 rpm using an eppendorf with a 0.45 μ m Durapore membrane (Millipore) and analyzed by HPLC.

HPLC Analysis. The concentration of the different products was analyzed by HPLC with a quaternary pump (Delta 600, Waters) coupled to a Lichrosorb-NH2 column (250 mm \times 4.6 mm) (Merck). The mobile phase was acetonitrile:water (75:25 v/v), conditioned with helium, and used at a flow rate of 0.7 mL/min. The column temperature was kept constant at 25 $^{\circ}$ C. A differential refractometric detector (model 9040, Varian) was used and set to a constant temperature of 30 $^{\circ}$ C. The data obtained were analyzed using the Millennium Software.

RESULTS AND DISCUSSION

Composition of Sugar Syrup and Molasses. Sugar syrup is the sugar juice obtained after beet juice evaporation, from which sucrose is crystallized. Molasses is the main byproduct of the sugar industry. Both sources are notably less expensive than pure sucrose for production of FOSs. The use of molasses for FOS synthesis has only been explored by Shin et al. in cultures of *A. pullulans* cells (9), whereas, to our knowledge, the sugar syrup has not been assayed as an alternative substrate in this process.

Sugar syrup and molasses are mixtures of sucrose and other carbohydrates. Molasses also has more unidentified components and some particulate materials (especially carbonates). **Table 1** shows the specifications of beet syrup and molasses that have

been used in this study. Sugar syrup and molasses were viscous solutions containing 620 and 570 mg/mL sucrose and 0.6 and 5.6% (w/w) of betaine (trimethyl glycine), respectively. The weight distribution of carbohydrates in syrup was 74.7% sucrose, 13.6% 1-kestose, 7.5% glucose, 3.5% nystose, and 0.7% fructose; molasses contained 95.7% sucrose, 2.9% glucose, and 1.4% of other carbohydrates. The presence of a considerable amount of FOS (17.1%) in the syrup is noteworthy. The 1-kestose content depends on the origin of the raw material and the type of technology adopted in beet processing (17, 18) and may cause shape and kinetic modifications of the final sucrose crystals. On the other hand, kestoses are becoming increasingly important due to their properties as prebiotic compounds. Other trisaccharides such as raffinose and neokestose were also present, especially in sugar syrup, but at a lower concentration (<0.2% w/w) as compared with 1-kestose.

Effect of pH and Substrate Concentration on FOS Production. The presence of a fructosyl transfer activity in commercial Pectinex Ultra SP-L from Novozymes A/S (used in juice clarification) was first reported by Hang and Woodams (14). Using Pectinex Ultra SP-L, we found that fructose was formed in a very small scale as compared with that of glucose, which indicated that this activity was better defined by a transfructosidase than a β -fructofuranosidase (16). Optimum pH and temperature for production of FOSs from sucrose using Pectinex Ultra SP-L were reported as 5.5 and 60 $^{\circ}$ C, respectively (19). Recently, we have purified the enzyme responsible for this transfructosidase activity (data not shown).

The initial pH of the sugar syrup and molasses was 7.5 and 8.9, respectively. We observed that FOS production from syrup increased from 30.6 to 36.4% when moving from the starting pH (7.5) to pH 5.5 (by the addition of glacial acetic acid), whereas for molasses an increment from 3.5 to 14.8% was observed when lowering the pH from 8.9 to 5.5.

To reduce the viscosity of the feedstock, which may cause technological difficulties, several dilutions from syrup and molasses were prepared by adding distilled water to the substrate and adjusting the pH to 5.5. This allowed us to investigate the effect of substrate concentration on FOS production.

The synthesis of FOSs from sucrose is a kinetically controlled reaction that involves a fructosyl-enzyme intermediate. The nucleophiles H₂O and sucrose compete for the fructosyl-enzyme intermediate. When H₂O is the nucleophile, the enzyme acts as a hydrolase (releasing glucose and fructose). When sucrose is the nucleophile, the enzyme acts as a transfructosidase. The first condensation product (1-kestose) can also be hydrolyzed by the enzyme. The reaction time must be carefully controlled to stop the process when the maximum yield of condensation products is achieved. The maximum yield of FOS depends on two parameters: the concentration of sucrose and the intrinsic transferase/hydrolase ratio of the enzyme. To improve the yield of transfructosylating products, a high sucrose concentration must be used to increase the ratio $k_3 \cdot [\text{sucrose}] / k_2 \cdot [\text{H}_2\text{O}]$ (20).

Different diluted solutions of syrup and molasses containing Pectinex Ultra SP-L (5 U/mL) were incubated at 60 $^{\circ}$ C for 24 h, and the FOS content was analyzed. **Table 2** summarizes the initial sucrose concentration, as well as the sucrose conversion and FOS production in 24 h. It is noteworthy that the FOS percentage (referred to the total amount of carbohydrates) maintained a value close to 53–57% (w/w) for syrup and 41–46% (w/w) for molasses, which confirmed that the transferase/hydrolase ratio for this enzyme must be considerably high. As expected, the maximum FOS production (460 mg/mL for syrup and 245 mg/mL for molasses) was obtained without dilution.

Table 2. Effect of Dilution of Sugar Syrup and Molasses on Production of FOS^a

substrate	dilution (mL H ₂ O added/ mL substrate)	sucrose consumed (%) ^b	1-kestose (%) ^c	nystose (%) ^c	1 ^F -fructosyl-nystose (%) ^c	total FOS (%) ^c	FOS production (mg/mL)
syrup	0.0	73	37.7	16.5	1.5	55.7	460
syrup	0.2	79	21.1	26.9	6.9	55.0	380
syrup	0.5	83	12.7	26.9	14.3	53.8	295
syrup	1.0	86	16.0	22.0	18.7	56.7	235
molasses	0.0	45	33.5	7.8	0.3	41.6	245
molasses	0.2	57	20.6	20.7	2.5	43.8	215
molasses	0.5	55	15.2	22.5	7.8	45.6	180
molasses	1.0	61	9.0	19.7	12.6	41.3	125

^a Experimental conditions: 5 U/mL fructosyltransferase, 60 °C, 220 rpm, and pH 5.5. Values refer to 24 h of reaction. ^b Percentage of the initial sucrose that is transformed into carbohydrates in 24 h. ^c The values refer to percentage of FOS with respect to the total amount of carbohydrates in the mixture, after 24 h of reaction. The initial percentages of 1-kestose and nystose in sugar syrup were 13.6 and 3.5%, respectively.







Figure 1. Time dependency of FOS batch production from (A) beet sugar syrup and (B) molasses, catalyzed by soluble Pectinex Ultra SP-L. Experimental conditions: pH 5.5, 5.0 U/m (standard DNS assay), and 60 °C. The substrate was diluted by adding 0.2 mL H₂O per mL.

In contrast, 1^F-fructosyl-nystose (GF₄) production increased with substrate dilutions (up to 12-fold with syrup and 42-fold with molasses, **Table 2**). In fact, at lower sucrose concentrations, the formation of high polymerization degree products is favored, as there exists more competition between sucrose and reaction products to accept a fructosyl unit.

Batch Production of FOSs. We analyzed the kinetics of FOS batch production of FOSs from sugar syrup and molasses. To have an efficient mixing, dilutions (1:1.2) from syrup and molasses were utilized. **Figure 1** shows the progress of the process, and **Figure 2** represents the percentage and concentra-

Figure 2. (A) Kinetics of FOS production and (B) percentage of FOS referred to the total amount of carbohydrates using sugar syrup (○) and molasses (●). Experimental conditions: pH 5.5, 5.0 U/mL (standard DNS assay), and 60 °C. The substrate was diluted by adding 0.2 mL H₂O per mL.

tion (mg/mL) of total FOS. In the case of syrup, the FOS concentration reached a maximum value of 388 mg/mL after 30 h (228 mg/mL 1-kestose, 149 mg/mL nystose, and 9 mg/mL 1^F-fructofuranosyl-nystose). At this reaction time, the percentage of FOS was 56.0%, referred to total carbohydrates in the mixture. For molasses, the maximum concentration of FOSs was about 235 mg/mL (49.2% FOS) after 65 h (62 mg/mL 1-kestose, 143 mg/mL nystose, and 30 mg/mL 1^F-fructofuranosyl-nystose). After 140 h, reactions were close to equilibrium, with a FOS percentages of 49 and 42% for syrup and




Figure 3. (A) FOS production and (B) percentage of FOS referred to the total amount of carbohydrates using sucrose (630 mg/mL), beet sugar syrup, and molasses, catalyzed by immobilized Pectinex Ultra SP-L in Sepabeads EC-EP5. Experimental conditions: 1.0 U/mL (standard DNS assay), pH 5.5, and 60 °C. The initial percentage of FOS in sugar syrup was 17.1% (142 mg/mL)—indicated by a black bar.

molasses, respectively (Figure 2). Similar results using pure sucrose as the substrate were also obtained in our laboratory with soluble and immobilized Pectinex Ultra SP-L (16).

Immobilization of Fructosyltransferase on Sepabeads EC-EP5. Sepabeads EC are polymethacrylate-based carriers for enzyme immobilization. The series Sepabeads EC-EP are epoxy-activated, with a high reactive group density. The porosity of these materials is very suitable to obtain biocatalysts with a high volumetric activity. The chemistry for attachment enzyme/support is straightforward. As compared with other epoxy acrylic polymers, Sepabeads EC-EP carriers possess high mechanical stability, low compressibility, and high resistance to microbial attack. Furthermore, the raw materials applied for the production of these supports are included in the EU list of resins allowed for the processing of foodstuffs (21).

Fructosyltransferase enzyme in Pectinex Ultra SP-L was immobilized on Sepabeads EC-EP5. Immobilization was carried out by mixing the Pectinex and the polymer, without any buffer or pH adjustment (16). The protein concentration in commercial preparation of the enzyme was 17.8 mg/mL. The activity toward sucrose, using the DNS assay, was 321 U/mL. The protein content of the immobilized biocatalyst was 36 mg/g of support and its activity 21.0 U/g biocatalyst.

Fructosyltransferase immobilized on Sepabeads FP-EC5 was utilized for FOS production. Pure sucrose (630 mg/mL), sugar syrup, and molasses were comparatively used as substrates. Figure 3 shows the FOS production and FOS percentage—referred to total carbohydrates in the mixture—with the different starting materials. It is interesting to note that these values

correlated well with those obtained with the soluble enzyme. The highest percentage of FOS in the final mixture (61%) was found with pure sucrose. This value is very close to that we obtained with the related support Sepabeads EC-EP3 (16) and comparable to the reported with other immobilized transfructosidases (22). The FOS percentage determined with sugar syrup and molasses (53 and 37.5%) can be considered very satisfactory. It is interesting to point out that, in terms of FOS production, the highest value was found with sugar syrup (440 mg/mL) as compared with 385 and 215 mg/mL obtained with pure sucrose and molasses, respectively. This is a simple consequence of the presence of FOS in the starting syrup.

As syrup and molasses are colored materials and the FOS specifications for human nutrition are very restrictive in this subject, the FOS obtained by the methods described in this work could be easily employed for animal feed. In recent years, some relevant changes have been introduced in the nutrition of farmed animals, to cover the needs of essential nutrients and/or to optimize feed utilization. Examples include the administration of immunostimulants or natural substances having an antibacterial effect. Probiotics are being already employed to control pathogenic bacteria, reduce fecal odor, and enhance growth performance. Research to date indicates positive effects of probiotics on health status and performance of companion animals, livestock, and poultry (23–25). The process for FOS synthesis described here is inexpensive, simple, efficient, and, employing the immobilized biocatalyst, easy to scale-up to different reactor configurations.

ACKNOWLEDGMENT

We are very grateful to Azucarera Ebro (Valladolid, Spain) for supplying beet sugar syrup and molasses. We thank Drs. Moreno Daminati and Paolo Caimi (Resindion S.R.L.) for providing us Sepabeads EC polymers and for technical help. We are grateful to Ramiro Martínez (Novozymes A/S, Spain) for providing us with Pectinex Ultra SP-L.

LITERATURE CITED

- Bornet, F. R. J.; Brouns, F.; Tashiro, Y.; Duvillier, V. Nutritional aspects of short-chain fructooligosaccharides: Natural occurrence, chemistry, physiology and health implications. *Dig. Liver Dis.* **2002**, *34*, S111–S120.
- Gibson, G. R.; Ottaway, R. A. *Prebiotics: New Developments in Functional Foods*; Chandos Publishing: Oxford, 2000.
- Tuohy, K. M.; Rouzard, G. C. M.; Bruck, W. M.; Gibson, G. R. Modulation of the human gut microflora towards improved health using prebiotics—Assessment of efficacy. *Curr. Pharm. Des.* **2005**, *11*, 75–90.
- Antosova, M.; Polakovic, M. Fructosyltransferases: The enzymes catalyzing production of fructooligosaccharides. *Chem. Pap.-Chem. Zvesti* **2001**, *55*, 350–358.
- Marx, S. P.; Winkler, S.; Hartmeier, W. Metabolization of beta-(2,6)-linked fructose-oligosaccharides by different bifidobacteria. *FEMS Microbiol. Lett.* **2000**, *182*, 163–169.
- Grizard, D.; Barthomeuf, C. Enzymatic synthesis and structure determination of NEO-FOS. *Food Biotechnol.* **1999**, *13*, 93–105.
- Hirayama, M.; Sumi, N.; Hidaka, H. Purification and properties of a fructooligosaccharide-producing beta-fructofuranosidase from *Aspergillus niger* ATCC-20611. *Agric. Biol. Chem.* **1989**, *53*, 667–673.
- Sangeetha, P. T.; Ramesh, M. N.; Prapulla, S. G. Fructooligosaccharide production using fructosyl transferase obtained from recycling culture of *Aspergillus oryzae* CFR 202. *Process Biochem.* **2005**, *40*, 1085–1088.

- (9) Shin, H. T.; Baig, S. Y.; Lee, S. W.; Suh, D. S.; Kwon, S. T.; Lim, Y. B.; Lee, J. H. Production of fructo-oligosaccharides from molasses by *Aureobasidium pullulans* cells. *Bioresour. Technol.* **2004**, *93*, 59–62.
- (10) Fernandez, R. C.; Maresma, B. G.; Juarez, A.; Martinez, J. Production of fructooligosaccharides by beta-fructofuranosidase from *Aspergillus* sp. 27H. *J. Chem. Technol. Biotechnol.* **2004**, *79*, 268–272.
- (11) Nguyen, Q. D.; Rezessy-Szabo, J. M.; Bhat, M. K.; Hoschke, A. Purification and some properties of beta-fructofuranosidase from *Aspergillus niger* IMI303386. *Process Biochem.* **2005**, *40*, 2461–2466.
- (12) Spiegel, J. E.; Rose, R.; Karabell, P.; Frankos, V. H.; Schmitt, D. F. Safety and benefits of fructooligosaccharides as food ingredients. *Food Technol.* **1994**, *48*, 85–89.
- (13) Okai, A. A. E.; Gierschner, K. Size and charge properties of the pectic and cellulolytic enzymes in a commercial enzyme preparation. *Z. Lebensm. Unters. Forsch.* **1991**, *192*, 244–248.
- (14) Hang, Y. D.; Woodams, E. E. Fructosyltransferase activity of commercial enzyme preparations used in fruit juice processing. *Biotechnol. Lett.* **1995**, *17*, 741–745.
- (15) Tanriseven, A.; Gokmen, F. Novel method for the production of a mixture containing fructooligosaccharides and isomaltoligosaccharides. *Biotechnol. Tech.* **1999**, *13*, 207–210.
- (16) Ghazi, I.; Gómez de Segura, A.; Fernández-Arrojo, L.; Alcalde, M.; Yates, M.; Rojas-Cervantes, M. L.; Plou, F. J.; Ballesteros, A. Immobilisation of fructosyltransferase from *Aspergillus aculeatus* on epoxy-activated Sepabeads EC for the synthesis of fructo-oligosaccharides. *J. Mol. Catal. B: Enzymol.* **2005**, *35*, 19–27.
- (17) Vaccari, G.; Lodi, G.; Tamburini, E.; Bernardi, T.; Tosi, S. Detection of oligosaccharides in sugar products using planar chromatography. *Food Chem.* **2001**, *74*, 99–110.
- (18) Vaccari, G.; Mantovani, G.; Sgualdino, G.; Tamburini, E.; Aquilano, D. Fructo-oligosaccharides and sucrose crystal growth morphology-I. Experimental growth habits. *Zuckerindustrie* **1999**, *124*, 34–39.
- (19) Tanriseven, A.; Aslan, Y. Immobilization of Pectinex Ultra SP-L to produce fructooligosaccharides. *Enzyme Microb. Technol.* **2005**, *36*, 550–554.
- (20) Buchholz, K.; Kasche, V.; Bornscheuer, U. T. *Biocatalysts and Enzyme Technology*; Wiley-VCH Verlag: Weinheim, 2005.
- (21) Resolution AP(97)1, On ion exchange and adsorbent resins used in the processing of foodstuffs (adopted by the EU Committee of Ministers on September 30, 1997).
- (22) Lee, W. C.; Chiang, C. J.; Tsai, P. Y. Kinetic modeling of fructo-oligosaccharide production catalyzed by immobilized beta-fructofuranosidase. *Ind. Eng. Chem. Res.* **1999**, *38*, 2564–2570.
- (23) Brambilla, G.; De Filippis, S. Trends in animal feed composition and the possible consequences on residue tests. *Anal. Chim. Acta* **2005**, *529*, 7–13.
- (24) Mosenthin, R.; Bauer, E. The potential use of prebiotics in pig nutrition. *Asian-Australas. J. Anim. Sci.* **2000**, *13*, 315–325.
- (25) Flickinger, E. A.; Fahey, G. C. Pet food and feed applications of inulin, oligofructose and other oligosaccharides. *Br. J. Nutr.* **2002**, *87*, S297–S300.

Received for review December 2, 2005. Revised manuscript received February 13, 2006. Accepted March 7, 2006. This research was supported by the Spanish Education and Science Ministry (Project BIO2004-03773-C04-01). We thank the Spanish AECL, Gobierno Vasco, and Instituto Danone for research fellowships.

JF053023B